

Access DB# 154101

## SEARCH REQUEST FORM

Scientific and Technical Information Center

Requester's Full Name: BEN SACLEY Examiner #: 73489 Date: 5/20/05  
Art Unit: 1626 Phone Number 302-20704 Serial Number: 10/609,088  
Mail Box and Bldg/Room Location: REM-5335 Results Format Preferred (circle): PAPER ☒ DISK ☐ E-MAIL ☐

If more than one search is submitted, please prioritize searches in order of need.

\*\*\*\*\*

Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: Process for Removal of Aroclor from Aqueous Solution  
Inventors (please provide full names): Ward et al.

Earliest Priority Filing Date: \_\_\_\_\_

\*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.

Removal of Aroclor from Aqueous Solution  
in the presence of an acid catalyst with a scavenger  
Compound containing a thiol or a hydroxyl group.

Best Available Copy

## STAFF USE ONLY

	Type of Search	Vendors and cost where applicable
Searcher: <u>K. Fuller</u>	NA Sequence (#) _____	STN <u>✓</u>
Searcher Phone #: _____	AA Sequence (#) _____	Dialog _____
Searcher Location: _____	Structure (#) <u>2</u>	Questel/Orbit _____
Date Searcher Picked Up: _____	Bibliographic _____	Dr.Link _____
Date Completed: <u>5/26/05</u>	Litigation _____	Lexis/Nexis _____
Searcher Prep & Review Time: <u>30</u>	Fulltext _____	Sequence Systems _____
Clerical Prep Time: _____	Patent Family _____	WWW/Internet _____
Online Time: <u>45</u>	Other _____	Other (specify) _____



# **STIC Search Report**

**EIC 1700**

SB31

STIC Database Tracking Number: 154101

**TO: Ben Sackey**  
**Location: REM 5B35**  
**Art Unit : 1626**  
**May 26, 2005**

**Case Serial Number: 10/609088**

**From: Kathleen Fuller**  
**Location: EIC 1700**  
**REMSSEN 4B28**  
**Phone: 571/272-2505**  
**Kathleen.Fuller@uspto.gov**

## **Search Notes**

=> FILE REG

FILE 'REGISTRY' ENTERED AT 15:21:57 ON 26 MAY 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 25 MAY 2005 HIGHEST RN 851163-60-5

DICTIONARY FILE UPDATES: 25 MAY 2005 HIGHEST RN 851163-60-5

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.

\*\*\*\*\*  
\*  
\* The CA roles and document type information have been removed from \*  
\* the IDE default display format and the ED field has been added, \*  
\* effective March 20, 2005. A new display format, IDERL, is now \*  
\* available and contains the CA role and document type information. \*  
\*  
\*\*\*\*\*

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=> FILE HCAPLUS

FILE 'HCAPLUS' ENTERED AT 15:22:02 ON 26 MAY 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS).

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 26 May 2005 VOL 142 ISS 22

FILE LAST UPDATED: 25 May 2005 (20050525/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=&gt; D QUE

L3 1 SEA FILE=REGISTRY ABB=ON ACROLEIN/CN  
 L4 1 SEA FILE=REGISTRY ABB=ON ACRYLONITRILE/CN  
 L11 12231 SEA FILE=HCAPLUS ABB=ON L3  
 L12 27790 SEA FILE=HCAPLUS ABB=ON L4  
 L13 3860 SEA FILE=HCAPLUS ABB=ON L12 (L) (PREP OR PUR)/RL  
 L14 1179 SEA FILE=HCAPLUS ABB=ON L11 AND L12  
 L15 200 SEA FILE=HCAPLUS ABB=ON L11 (L) REM/RL  
 L16 14 SEA FILE=HCAPLUS ABB=ON L13 AND L15  
 L17 2 SEA FILE=REGISTRY ABB=ON 64-19-7 OR 79-14-1  
 L18 98066 SEA FILE=HCAPLUS ABB=ON L17  
 L19 2276 SEA FILE=HCAPLUS ABB=ON L18 (L) CAT/RL  
 L20 3 SEA FILE=HCAPLUS ABB=ON L14 AND L19  
 L21 20 SEA FILE=HCAPLUS ABB=ON L14 AND ACID (L) CAT/RL  
 L22 4 SEA FILE=REGISTRY ABB=ON 60-23-1 OR 60-24-2 OR 68-11-1 OR  
 123-81-9  
 L23 18871 SEA FILE=HCAPLUS ABB=ON L22  
 L24 18 SEA FILE=HCAPLUS ABB=ON L14 AND L23  
 L25 755 SEA FILE=HCAPLUS ABB=ON ACROLEIN (L) REMOV?  
 L26 157 SEA FILE=HCAPLUS ABB=ON ACROLEIN (4A) REMOV?  
 L27 1 SEA FILE=HCAPLUS ABB=ON L24 AND L26  
 L28 504 SEA FILE=HCAPLUS ABB=ON L22/DP  
 L29 3 SEA FILE=HCAPLUS ABB=ON L24 AND L28  
 L30 35 SEA FILE=HCAPLUS ABB=ON L16 OR L20 OR L21 OR L27 OR L29  
 L31 15 SEA FILE=HCAPLUS ABB=ON L24 AND (SCAVENG? OR ?THIOL? OR  
 HYDROXY? OR OH)  
 L32 1 SEA FILE=HCAPLUS ABB=ON (L25 OR L15) AND L31  
 L33 35 SEA FILE=HCAPLUS ABB=ON L30 OR L32.

=&gt; D L33.BIB ABS HITIND HITSTR 1-35

L33 ANSWER 1 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2005:2238 HCAPLUS  
 DN 142:94278  
 TI Process for the removal of acrolein from acrylonitrile  
 product streams via acetal formation and distillation  
 IN Ward, Gregory J.; Blanchard, Bryan C.; Moffatt, Scott G.; Monical, Valerie  
 S.; Murphy, Richard D.; Ramchandran, Balshekar  
 PA USA  
 SO U.S. Pat. Appl. Publ., 5 pp.  
 CODEN: USXXCO  
 DT Patent  
 LA English  
 FAN.CNT 1

*applicant*

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2004267054	A1	20041230	US 2003-609088	20030627
	WO 2005003063	A2	20050113	WO 2004-US19826	20040621
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRAI US 2003-609088 A 20030627

AB **Acrolein** is removed from a process stream such as a process stream generated in the manufacture of acrylonitrile. The process includes reacting the **acrolein** with a compound containing a reactable **thiol** or **hydroxyl** moiety (e.g., 2-mercaptoethanol) in the presence of an acid catalyst (e.g., acetic acid) followed by distillation. These refined process streams contain  $\leq 5$  ppm unreacted **acrolein**.

IC ICM C07C043-32

INCL 568594000

CC 35-2 (Chemistry of Synthetic High Polymers)

Section cross-reference(s): 23, 48

IT Acids, uses

RL: CAT (Catalyst use); USES (Uses)

(acetalization catalysts in a process for the removal of **acrolein** from acrylonitrile product streams via acetal formation and distillation)

IT Alcohols, reactions

**Thiols** (organic), reactions

RL: RGT (Reagent); RACT (Reactant or reagent)

(acetals with **acrolein**; in a process for the removal of **acrolein** from acrylonitrile product streams via acetal formation and distillation)

IT Acetalization catalysts

(acids; in a process for the removal of **acrolein**

from acrylonitrile product streams via acetal formation and distillation)

IT Acetalization

(in a process for the removal of **acrolein** from

acrylonitrile product streams via acetal formation and distillation)

IT 64-19-7, Acetic acid, uses 79-14-1, Glycolic acid, uses

RL: CAT (Catalyst use); USES (Uses)

(acetalization catalysts in a process for the removal of **acrolein** from acrylonitrile product streams via acetal formation and distillation)

IT 60-23-1DP, 1-Amino-2-mercaptoethane, acetals with **acrolein**

60-24-2DP, 2-Mercaptoethanol, acetals with **acrolein**

68-11-1DP, Mercaptoacetic acid, acetals with **acrolein**

123-81-9DP, Ethylene glycol bithioglycolate, acetals with **acrolein**

RL: BYP (Byproduct); REM (Removal or disposal); PREP (Preparation); PROC (Process)

(in a process for the removal of **acrolein** from

acrylonitrile product streams via acetal formation and distillation)

IT 7732-18-5, Water, uses

RL: NUU (Other use, unclassified); USES (Uses)

(in a process for the removal of **acrolein** from

acrylonitrile product streams via acetal formation and distillation)

IT 107-02-8DP, Acrolein, acetals with alcs. or thiols

RL: BYP (Byproduct); REM (Removal or disposal); PREP

(Preparation); PROC (Process)

(process for the removal of **acrolein** from

acrylonitrile product streams via acetal formation and distillation)

IT 107-13-1P, Acrylonitrile, preparation

RL: PEP (Physical, engineering or chemical process); PUR

(Purification or recovery); PYP (Physical process); PREP


(Preparation); PROC (Process)

(process for the removal of **acrolein** from


acrylonitrile product streams via acetal formation and distillation)

IT 64-19-7, Acetic acid, uses 79-14-1, Glycolic acid, uses  
 RL: CAT (Catalyst use); USES (Uses)  
 (acetalization catalysts in a process for the removal of acrolein from acrylonitrile product streams via acetal formation and distillation)

RN 64-19-7 HCAPLUS  
 CN Acetic acid (7CI, 8CI, 9CI) (CA INDEX NAME)




RN 79-14-1 HCAPLUS  
 CN Acetic acid, hydroxy- (9CI) (CA INDEX NAME)



IT 60-23-1DP, 1-Amino-2-mercaptoethane, acetals with acrolein  
 60-24-2DP, 2-Mercaptoethanol, acetals with acrolein  
 68-11-1DP, Mercaptoacetic acid, acetals with acrolein  
 123-81-9DP, Ethylene glycol bithioglycolate, acetals with acrolein  
 RL: BYP (Byproduct); REM (Removal or disposal); PREP (Preparation); PROC (Process)  
 (in a process for the removal of acrolein from acrylonitrile product streams via acetal formation and distillation)


RN 60-23-1 HCAPLUS  
 CN Ethanethiol, 2-amino- (8CI, 9CI) (CA INDEX NAME)




RN 60-24-2 HCAPLUS  
 CN Ethanol, 2-mercapto- (8CI, 9CI) (CA INDEX NAME)




RN 68-11-1 HCAPLUS  
 CN Acetic acid, mercapto- (8CI, 9CI) (CA INDEX NAME)



RN 123-81-9 HCAPLUS  
 CN Acetic acid, mercapto-, 1,2-ethanediyl ester (9CI) (CA INDEX NAME)



IT 107-02-8DP, Acrolein, acetals with alcs. or  
thiols  
RL: BYP (Byproduct); REM (Removal or disposal); PREP  
(Preparation); PROC (Process)  
(process for the removal of acrolein from  
acrylonitrile product streams via acetal formation and distillation)  
RN 107-02-8 HCAPLUS  
CN 2-Propenal (9CI) (CA INDEX NAME)



IT 107-13-1P, Acrylonitrile, preparation  
RL: PEP (Physical, engineering or chemical process); PUR  
(Purification or recovery); PYP (Physical process); PREP  
(Preparation); PROC (Process)  
(process for the removal of acrolein from  
acrylonitrile product streams via acetal formation and distillation)  
RN 107-13-1 HCAPLUS  
CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 2 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
AN 2004:856907 HCAPLUS  
DN 141:356031  
TI Functionalized nanotubes  
IN Fischer, Alan; Hoch, Robert; Moy, David; Lu, Ming; Martin, Mark; Niu, Chun  
Ming; Ogata, Naoya; Tennent, Howard; Dong, Liwen; Sun, Ji; Helms, Larry;  
Jameison, Fabian; Liang, Pam; Simpson, David  
PA Hyperion Catalysis International, Inc., USA  
SO U.S. Pat. Appl. Publ., 50 pp., Cont.-in-part of U.S. Ser. No. -594,673.  
CODEN: USXXCO  
DT Patent  
LA English  
FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 2004202603	A1	20041014	US 2004-837125	20040430
	US 6203814	B1	20010320	US 1994-352400	19941208
PRAI	US 1994-352400	A3	19941208		
	US 1996-611368	B1	19960306		
	US 1996-37238P	P	19960925		
	US 1997-812856	B1	19970306		
	US 2000-594673	A2	20000616		

AB The invention describes graphitic nanotubes, which includes tubular fullerenes (commonly called "buckytubes") and fibrils, which are functionalized by chemical substitution or by adsorption of functional moieties. More specifically the invention relates to graphitic nanotubes which are uniformly or non-uniformly substituted with chemical moieties or upon which certain cyclic compds. are adsorbed and to complex structures

comprised of such functionalized nanotubes linked to one another. The invention also relates to methods for introducing functional groups onto the surface of such nanotubes. The invention further relates to uses for functionalized nanotubes.

IC ICM D01F009-12

ICS C07C063-333

INCL 423447200; 562492000; 564426000

CC 66-4 (Surface Chemistry and Colloids)

Section cross-reference(s): 7


IT 56-87-1DP, L-Lysine, carbon fibril bonded, preparation 58-85-5DP, Biotin, surface reaction product with carbon fibrils 60-24-2DP, Monothioethylene glycol, surface reaction product with carbon nanotubes and fibrils 75-89-8DP, 2,2,2-Trifluoroethanol, surface reaction product with carbon nanotubes and fibrils 79-06-1DP, 2-Propenamide, surface reaction product with carbon nanotubes and fibrils 79-10-7DP, 2-Propenoic acid, surface reaction product with carbon nanotubes and fibrils 107-02-8DP, Propenal, surface reaction product with carbon nanotubes and fibrils 107-11-9DP, 3-Amino-1-propene, surface reaction product with carbon nanotubes and fibrils 107-13-1DP, 2-Propenenitrile, surface reaction product with carbon nanotubes and fibrils 107-18-6DP, 2-Propen-1-ol, surface reaction product with carbon nanotubes and fibrils 108-31-6DP, 2,5-Furandione, surface reaction product with carbon nanotubes and fibrils 109-72-8DP, Butyllithium, surface reaction product with carbon nanotubes and fibrils 110-16-7DP, 2-Butenedioic acid (Z)-, surface reaction product with carbon nanotubes and fibrils 111-86-4DP, 1-Octanamine, surface reaction product with carbon nanotubes and fibrils 124-30-1DP, 1-Octadecanamine, surface reaction product with carbon nanotubes and fibrils 151-50-8DP, Potassium cyanide, surface reaction product with carbon nanotubes and fibrils 530-62-1DP, N,N'-Carbonyl diimidazole, surface reaction product with carbon nanotubes and fibrils 593-56-6DP, Methoxyamine hydrochloride, surface reaction product with carbon nanotubes and fibrils 814-68-6DP, Propenoyl chloride, surface reaction product with carbon nanotubes and fibrils 994-30-9DP, Chlorotriethylsilane, surface reaction product with carbon nanotubes and fibrils 1310-73-2DP, Sodium hydroxide, surface reaction product with carbon nanotubes and fibrils 1333-74-0DP, Hydrogen, surface reaction product with carbon nanotubes and fibrils 1336-21-6DP, Ammonium hydroxide, surface reaction product with carbon nanotubes and fibrils 1892-57-5DP, 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide, surface reaction product with carbon nanotubes and fibrils 2016-57-1DP, 1-Aminodecane, surface reaction product with carbon nanotubes and fibrils 2074-87-5DP, Cyanogen, surface reaction product with carbon nanotubes and fibrils 4048-33-3DP, 6-Aminohexan-1-ol, surface reaction product with carbon nanotubes and fibrils 4781-83-3DP, 2-Iminothiolane hydrochloride, surface reaction product with carbon nanotubes and fibrils 5591-94-6DP, surface reaction product with carbon nanotubes and fibrils 5957-17-5DP, Triethyl(2-hydroxyethyl)ammonium iodide, surface reaction product with carbon nanotubes and fibrils 7664-41-7DP, Ammonia, surface reaction product with carbon nanotubes and fibrils 7664-93-9DP, Sulfuric acid, surface reaction product with carbon nanotubes and fibrils 7697-37-2DP, Nitric acid, surface reaction product with carbon nanotubes and fibrils 7704-34-9DP, Sulfur, surface reaction product with carbon nanotubes and fibrils 7732-18-5DP, Water, surface reaction product with carbon nanotubes and fibrils 7782-44-7DP, Oxygen, surface reaction product with carbon nanotubes and fibrils 13214-66-9DP, 4-Phenylbutylamine, surface reaction product with carbon nanotubes and fibrils 19008-71-0DP, 8-Aminooctan-1-ol, surface reaction product with carbon nanotubes and fibrils 23160-46-5DP, 10-Aminodecan-1-ol, surface reaction product with carbon nanotubes and fibrils 103708-09-4DP, Sulfosuccinimidyl-4-(N-




maleimidomethyl)cyclohexanecarboxylate, surface reaction product with carbon nanotubes and fibrils 142755-63-3DP, 18-Aminooctadecan-1-ol, surface reaction product with carbon nanotubes and fibrils  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (surface functionalization of carbon nanotubes and fibrils for enzyme immobilization)

IT 60-24-2DP, Monothioethylene glycol, surface reaction product with carbon nanotubes and fibrils 107-02-8DP, Propenal, surface reaction product with carbon nanotubes and fibrils 107-13-1DP, 2-Propenenitrile, surface reaction product with carbon nanotubes and fibrils  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (surface functionalization of carbon nanotubes and fibrils for enzyme immobilization)

RN 60-24-2 HCAPLUS  
 CN Ethanol, 2-mercapto- (8CI, 9CI) (CA INDEX NAME)



RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 3 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2004:848394 HCAPLUS  
 DN 142:37985  
 TI Palladium-catalyzed Mizoroki-Heck-type reactions using telluronium salts  
 AU Hirabayashi, Kazunori; Nara, Yoshiko; Shimizu, Toshio; Kamigata, Nobumasa  
 CS Department of Chemistry, Graduate School of Science, Tokyo Metropolitan University, Tokyo, 192-0397, Japan  
 SO Chemistry Letters (2004), 33(10), 1280-1281  
 CODEN: CMLTAG; ISSN: 0366-7022  
 PB Chemical Society of Japan  
 DT Journal  
 LA English  
 AB A Mizoroki-Heck-type reaction of telluronium iodides with olefins proceeded under mild conditions to produce substituted olefins in high yields. The reaction required a catalytic amount of palladium(II) species and a stoichiometric amount of silver(I) acetate as an additive. For example, the acetic acid ester palladium(2+) salt/acetic acid ester silver(1+) salt-catalyzed phenylation of 2-propenoic acid ester Bu ester using dimethyl(phenyl)telluronium iodide gave (2E)-3-phenyl-2-propenoic acid Bu ester in 99% yield.  
 CC 25-18 (Benzene, Its Derivatives, and Condensed Benzenoid Compounds)  
 Section cross-reference(s): 29  
 IT 107-02-8, 2-Propenal, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of (phenyl)propenal by Mizoroki-Heck reaction using dimethyl(phenyl)telluronium iodide and acrylaldehyde as starting materials and palladium as catalyst and silver compound as additive)

IT 107-13-1, 2-Propenenitrile, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation of (phenyl)propenenitrile by Mizoroki-Heck reaction using dimethyl(phenyl)telluronium iodide and acrylonitrile as starting materials and palladium as catalyst and silver compound as additive)


IT 3375-31-3  
 RL: CAT (Catalyst use); USES (Uses)  
 (preparation of (phenyl)propenoic acid ester by Mizoroki-Heck reaction using dimethyl(phenyl)telluronium iodide and Bu acrylate as starting materials and palladium acetate as catalyst and silver compound as additive)

IT 563-63-3  
 RL: CAT (Catalyst use); USES (Uses)  
 (preparation of (phenyl)propenoic acid ester by Mizoroki-Heck reaction using dimethyl(phenyl)telluronium iodide and Bu acrylate as starting materials and palladium as catalyst and silver acetate as additive)

IT 7647-10-1, Palladium chloride (PdCl<sub>2</sub>) 13965-03-2, Dichlorobis(triphenylphosphine)palladium  
 RL: CAT (Catalyst use); USES (Uses)  
 (preparation of (phenyl)propenoic acid ester by Mizoroki-Heck reaction using dimethyl(phenyl)telluronium iodide and Bu acrylate as starting materials and palladium as catalyst and silver compound as additive)

IT 107-02-8, 2-Propenal, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation of (phenyl)propenal by Mizoroki-Heck reaction using dimethyl(phenyl)telluronium iodide and acrylaldehyde as starting materials and palladium as catalyst and silver compound as additive)

RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



IT 107-13-1, 2-Propenenitrile, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation of (phenyl)propenenitrile by Mizoroki-Heck reaction using dimethyl(phenyl)telluronium iodide and acrylonitrile as starting materials and palladium as catalyst and silver compound as additive)

RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 4 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2002:469853 HCAPLUS  
 DN 137:154608  
 TI Quantitative Characterization of the Local Electrophilicity of Organic Molecules. Understanding the Regioselectivity on Diels-Alder Reactions  
 AU Domingo, Luis R.; Aurell, M. Jose; Perez, Patricia; Contreras, Renato

CS Instituto de Ciencia Molecular Departamento de Quimica Organica,  
 Universidad de Valencia, Valencia, 46100, Spain

SO Journal of Physical Chemistry A (2002), 106(29), 6871-6875  
 CODEN: JPCAFH; ISSN: 1089-5639

PB American Chemical Society

DT Journal

LA English

AB Regional electrophilicity at the active sites of the reagents involved in polar Diels-Alder processes may be described on a quant. basis using an extension of the global electrophilicity index recently introduced by Parr et al. (J. Am. Chemical Society 1999, 121, 1922). The local or regional electrophilicity provides useful clues about the expected regioselectivity of the products on Diels-Alder reactions showing significant polar character. The local (regional) electrophilicity index shows significant advantages over other unnormalized definitions of relative electrophilicity proposed in the literature in the sense that it clearly identifies the relevant electrophilic sites in a mol. without restricting the search to those sites having comparable values of regional electrophilic/nucleophilic softness.

CC 22-5 (Physical Organic Chemistry)


IT 185313-96-6  
 RL: CAT (Catalyst use); CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent); USES (Uses)  
 (Lewis acid catalyst model; quant. characterization of the local electrophilicity of organic mols. and its use as a clue about the expected regioselectivity of Diels-Alder reactions)

IT 74-85-1, Ethylene, reactions 74-86-2, Acetylene, reactions 78-79-5, 2-Methyl-1,3-butadiene, reactions 78-94-4, Methyl vinyl ketone, reactions 96-33-3, Methyl acrylate 106-99-0, 1,3-Butadiene, reactions 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile, reactions 107-25-5, Methyl vinyl ether 922-64-5, 1,1-Dicyanoethylene 926-56-7, 4-Methyl-1,3-pentadiene 1515-77-1, 1-(Dimethylamino)-1,3-butadiene 2004-70-8, trans-1,3-Pentadiene 3036-66-6, 1-Methoxy-1,3-butadiene 3638-64-0, Nitroethylene 5763-87-1, (Dimethylamino)ethylene 6651-43-0, 1-(Trimethylsilyloxy)-1,3-butadiene 38053-91-7, 2-(Trimethylsilyloxy)-1,3-butadiene 51943-18-1, Methanamine, N-methylene-, conjugate acid  
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent)  
 (quant. characterization of the local electrophilicity of organic mols. and its use as a clue about the expected regioselectivity of Diels-Alder reactions)

IT 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile, reactions  
 RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent)  
 (quant. characterization of the local electrophilicity of organic mols. and its use as a clue about the expected regioselectivity of Diels-Alder reactions)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 5 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2002:119267 HCAPLUS

DN 136:167799

TI Process and apparatus for removing organic substances from a gas mixture  
IN Prinz, Peter; Schummer, Gunter; Schmitz, Josef; Strobel, Rainer; Stuwe, Arnd

PA Ec Erdoelchemie GmbH, Germany; Bayer Ag

SO Eur. Pat. Appl., 8 pp.

CODEN: EPXXDW

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1179359	A2	20020213	EP 2001-306341	20010724
	EP 1179359	A3	20020619		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	DE 10037774	A1	20020214	DE 2000-10037774	20000803
	ZA 2001006183	A	20030127	ZA 2001-6183	20010726
	US 2002087023	A1	20020704	US 2001-916336	20010730
	US 6541652	B2	20030401		
	CA 2354741	AA	20020203	CA 2001-2354741	20010802
	CN 1341471	A	20020327	CN 2001-132577	20010803
	JP 2002121180	A2	20020423	JP 2001-237174	20010803
	BG 105777	A	20020430	BG 2001-105777	20010803
PRAI	DE 2000-10037774	A	20000803		

AB Process for isolating one or more organic substances from a gas mixture in which these organic substances are present, in which the gas mixture is subjected to a quenching in a column, characterized in that quenching is carried out in the upper part of the column and the quenching liquid is subjected to stripping in the lower part of the column. The method is suitable for use in the manufacture of acrylonitrile or methacrylonitrile.

IC ICM B01D053-00

ICS B01D053-14; B01D053-18

CC 35-2 (Chemistry of Synthetic High Polymers)

IT 64-19-7, Acetic acid, processes 74-90-8, Hydrogen cyanide, processes

75-05-8, Acetonitrile, processes 79-10-7, Acrylic acid, processes

100-54-9, Nicotinonitrile 107-02-8, Acrolein, processes

124-38-9, Carbon dioxide, processes 288-42-6, Oxazole 630-08-0, Carbon

monoxide, processes 764-42-1, Fumaronitrile 7727-37-9, Nitrogen,

processes 7782-44-7, Oxygen, processes 14798-03-9, Ammonium, processes

RL: FMU (Formation, unclassified); REM (Removal or disposal);

FORM (Formation, nonpreparative); PROC (Process)

(process and apparatus for removing organic substances from a gas mixture

such as

acrylonitrile synthesis gases)

IT 107-13-1P, Acrylonitrile, preparation 126-98-7P,


Methacrylonitrile

RL: IMF (Industrial manufacture); PREP (Preparation)

(process and apparatus for removing organic substances from a gas mixture

such as

acrylonitrile synthesis gases)  
 IT 107-02-8, Acrolein, processes  
 RL: FMU (Formation, unclassified); REM (Removal or disposal);  
 FORM (Formation, nonpreparative); PROC (Process)  
 (process and apparatus for removing organic substances from a gas mixture  
 such as  
 acrylonitrile synthesis gases)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



IT 107-13-1P, Acrylonitrile, preparation  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (process and apparatus for removing organic substances from a gas mixture  
 such as  
 acrylonitrile synthesis gases)  
 RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 6 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2001:895585 HCAPLUS  
 DN 136:38504  
 TI Rosin-fatty acid vinylic polymers as moisture vapor barrier coatings  
 IN Shah, Rajnikant; Adams, Stanley C.  
 PA Westvaco Corporation, USA  
 SO U.S., 8 pp., Cont.-in-part of U.S. Ser. No. 174,398, abandoned.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 FAN.CNT 6

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6329068	B1	20011211	US 2000-545668	20000407
PRAI	US 1998-174398	B2	19981014		

AB This invention relates to novel rosin-fatty acid vinylic polymer compns.  
 which exhibit properties that make them useful for formulating moisture  
 vapor barrier coatings for paper and other cellulosic base materials. In  
 particular, this invention relates to rosin-fatty acid vinylic polymer  
 moisture barrier coatings obtained from the addition polymerization reaction  
 of a  
 mixture of rosin and fatty acid and a mixture of (meth)acrylic and vinylic  
 monomers.  
 IC ICM B32B023-08  
 INCL 428514000  
 CC 38-3 (Plastics Fabrication and Uses)  
 Section cross-reference(s): 42, 43  
 IT 78-67-1, AIBN 80-15-9, Cumene hydroperoxide 94-36-0, Benzoyl peroxide,  
 uses 110-05-4, tert-Butyl peroxide 614-45-9, tert-Butyl peroxybenzoate  
 13467-82-8, tert-Butyl peroctoate  
 RL: CAT (Catalyst use); USES (Uses)  
 (catalyst; rosin-fatty acid vinylic polymers as moisture  
 vapor barrier coatings)

IT. 78-85-3DP, Methacrolein, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 79-06-1DP, Acrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 79-10-7DP, Acrylic acid, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 79-39-0DP, Methacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 79-41-4DP, Methacrylic acid, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 80-62-6DP, Methyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 80-63-7DP, Methyl  $\alpha$ -chloroacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 96-05-9DP, Allyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 96-33-3DP, Methyl acrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 97-63-2DP, Ethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 97-86-9DP, Isobutyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 97-88-1DP, Butyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 98-83-9DP,  $\alpha$ -Methylstyrene, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 100-42-5DP, Styrene, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 101-43-9DP, Cyclohexyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 105-16-8DP, N,N-Diethylaminoethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 106-91-2DP, Glycidyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 107-02-8DP, Acrolein, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 107-13-1DP, Acrylonitrile, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 126-98-7DP, Methacrylonitrile, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 137-05-3DP, Methyl 2-cyanoacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 140-88-5DP, Ethyl acrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 141-32-2DP, Butyl acrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 142-09-6DP, Hexyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 142-90-5DP, Lauryl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 585-07-9DP, tert-Butyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 688-84-6DP, 2-Ethylhexyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 689-12-3DP, Isopropyl acrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 816-74-0DP, Methallyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 868-77-9DP, 2-Hydroxyethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 923-26-2DP, 2-Hydroxypropyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 925-60-0DP, Propyl acrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 1611-83-2DP, N-Phenylmethacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 1888-94-4DP, 2-Chloroethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 2039-87-4DP, o-Chlorostyrene, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 2156-96-9DP, Decyl acrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and viny compds. 2157-01-9DP, n-Octyl methacrylate, polymers with rosin

fatty acids and (meth)acrylic acid or esters and vinyl compds.  
 2177-42-6DP, 2-Nitro-2-methylpropyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds.  
 2177-70-0DP, Phenyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2210-28-8DP, Propyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2370-63-0DP, 2-Ethoxyethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2455-24-5DP, Tetrahydrofurfuryl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2495-37-6DP, Benzyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2675-94-7DP, N,N-Diethylacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2849-98-1DP, Amyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2867-47-2DP, N,N-Dimethylaminoethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 2998-18-7DP, sec-Butyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 3063-94-3DP, Hexafluoroisopropyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 3454-28-2DP, Furfuryl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 3683-12-3DP, 2-Phenylethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 3775-90-4DP, tert-Butylaminoethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 3887-02-3DP, N-Methylmethacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 4655-34-9DP, Isopropyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 5138-86-3DP, 2-Ethylbutyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 5441-99-6DP, N,N-Diethylmethacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 5883-17-0DP, N-Ethylacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 6976-91-6DP, N,N-Dimethylmethacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 6976-96-1DP, 3-Methoxybutyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 7336-27-8DP, Isoamyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 7370-88-9DP, N-Ethylmethacrylamide, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 7376-45-6DP, Crotyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 10595-06-9DP, 2-Phenoxyethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 10595-80-9DP, 2-Sulfoethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 13532-94-0DP, 2-n-Butoxyethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 13861-22-8DP, Propargyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 16868-14-7DP, Cyclopentyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 25013-15-4DP, Vinyltoluene, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 25338-51-6DP, tert-Butylstyrene, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 31736-34-2DP, Cinnamyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 38785-10-3DP, Trifluoroethyl methacrylate, polymers with rosin fatty acids and (meth)acrylic acid or esters and vinyl compds. 52858-59-0DP, Tetrahydropyranyl methacrylate, polymers with rosin fatty acids and

(meth)acrylic acid or esters and vinyl compds. 67905-44-6DP,  
2-Methoxybutyl methacrylate, polymers with rosin fatty acids and  
(meth)acrylic acid or esters and vinyl compds.

RL: IMF (Industrial manufacture); PEP (Physical, engineering or chemical  
process); PYP (Physical process); TEM (Technical or engineered material  
use); PREP (Preparation); PROC (Process); USES (Uses)

(rosin-fatty acid vinyl polymers as moisture vapor barrier coatings)


IT 107-02-8DP, Acrolein, polymers with rosin fatty acids and  
(meth)acrylic acid or esters and vinyl compds. 107-13-1DP,  
Acrylonitrile, polymers with rosin fatty acids and (meth)acrylic acid or  
esters and vinyl compds.

RL: IMF (Industrial manufacture); PEP (Physical, engineering or chemical  
process); PYP (Physical process); TEM (Technical or engineered material  
use); PREP (Preparation); PROC (Process); USES (Uses)

(rosin-fatty acid vinyl polymers as moisture vapor barrier coatings)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 7 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2001:391919 HCAPLUS

DN 134:371325

TI Oxidation and ammoxidation of acrylonitrile process waste water organics

IN Cesa, Mark Clark; Graham, Anne Marie; Shuki, Albert Richard

PA The Standard Oil Company, USA

SO U.S., 7 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6238574	B1	20010529	US 1998-182774	19981030
PRAI	US 1998-182774		19981030		

AB A process for upgrading aqueous acrylonitrile waste streams containing organic  
material comprising atomizing an acrylonitrile wastewater stream containing  
organic material, introducing the atomized acrylonitrile wastewater stream at  
a temperature below the decomposition temperature of the orgs. present in the  
wastewater

stream, into a reaction zone containing a catalyst and at least one reactant  
gas, reacting the atomized wastewater stream and reactant gas in the  
presence of the catalyst to convert at least some of the orgs. in the  
wastewater stream into at least one compound selected from the group  
consisting of acetonitrile, hydrogen cyanide and acrylonitrile.

IC ICM C02F001-72

INCL 210763000

CC 60-2 (Waste Treatment and Disposal)



Section cross-reference(s): 35

IT 107-13-1P, Acrylonitrile, processes  
 RL: FMU (Formation, unclassified); IMF (Industrial manufacture); PEP (Physical, engineering or chemical process); FORM (Formation, nonpreparative); **PREP (Preparation)**; PROC (Process)  
 (oxidation and ammoxidn. of acrylonitrile process wastewater orgs. to recover acetonitrile, hydrogen cyanide or acrylonitrile)

IT 79-06-1, Acrylamide, processes 107-02-8, Acrolein, processes 110-61-2, Succinonitrile 110-86-1, Pyridine, processes 288-13-1, Pyrazole 764-42-1, Fumaronitrile 1656-48-0, Bis(2-cyanoethyl)ether  
 RL: PEP (Physical, engineering or chemical process); **REM (Removal or disposal)**; PROC (Process)  
 (oxidation and ammoxidn. of acrylonitrile process wastewater orgs. to recover acetonitrile, hydrogen cyanide or acrylonitrile)


IT 107-13-1P, Acrylonitrile, processes  
 RL: FMU (Formation, unclassified); IMF (Industrial manufacture); PEP (Physical, engineering or chemical process); FORM (Formation, nonpreparative); **PREP (Preparation)**; PROC (Process)  
 (oxidation and ammoxidn. of acrylonitrile process wastewater orgs. to recover acetonitrile, hydrogen cyanide or acrylonitrile)

RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



IT 107-02-8, Acrolein, processes  
 RL: PEP (Physical, engineering or chemical process); **REM (Removal or disposal)**; PROC (Process)  
 (oxidation and ammoxidn. of acrylonitrile process wastewater orgs. to recover acetonitrile, hydrogen cyanide or acrylonitrile)

RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 8 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2001:152276 HCAPLUS  
 DN 134:193859  
 TI Polyoxometallate supported catalysts for conversion of alkanes  
 IN Devlin, Anna Marie; Volpe, Anthony Frank, Jr.  
 PA Rohm and Haas Company, USA  
 SO Eur. Pat. Appl., 19 pp.  
 CODEN: EPXXDW  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1078687	A1	20010228	EP 2000-306863	20000810
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	US 6387841	B1	20020514	US 2000-626640	20000727
	CA 2315884	AA	20010223	CA 2000-2315884	20000814

BR 2000003694	A	20010327	BR 2000-3694	20000818
CN 1285240	A	20010228	CN 2000-124283	20000823
CN 1129475	B	20031203		
JP 2001087653	A2	20010403	JP 2000-252555	20000823
US 2002142914	A1	20021003	US 2002-100946	20020319
PRAI US 1999-150309P	P	19990823		
US 2000-626640	A3	20000727		

AB These title catalysts are for example mixed metal oxides, optionally heteropolyacids on a polyoxometallate and used for the conversion of alkanes to unsatd. organic compds. Thus, propane was converted to acrylic acid at 380° in 3 s using a calcined 0.24 g Mo1V0.03Te0.23Nb0.08Ox catalyst on 1 g Cs3PMo12O40, showing selectivity 27.8 and acrylic acid yield 1.0%.

IC ICM B01J027-188  
ICS B01J027-198; C07C051-215; C07C005-42


CC 35-2 (Chemistry of Synthetic High Polymers)  
Section cross-reference(s): 67

IT 78-85-3P, Methacrolein 79-10-7P, Acrylic acid, preparation 79-41-4P, Methacrylic acid, preparation 106-99-0P, Butadiene, preparation 107-02-8P, Acrolein, preparation 107-13-1P, Acrylonitrile, preparation 108-31-6P, Maleic anhydride, preparation 110-16-7P, Maleic acid, preparation 115-07-1P, Propylene, preparation 115-11-7P, Isobutylene, preparation 126-98-7P, Methacrylonitrile 25167-67-3P, Butylene  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(polyoxometallate supported mixed metal oxide catalysts for conversion of alkanes)

IT 11104-88-4, Phosphomolybdic acid 12026-91-4 68335-84-2  
93222-18-5 121072-30-8 133644-76-5 312492-53-8 326859-62-5  
RL: CAT (Catalyst use); USES (Uses)  
(support; polyoxometallate supported mixed metal oxide catalysts for conversion of alkanes)

IT 107-02-8P, Acrolein, preparation 107-13-1P, Acrylonitrile, preparation  
RL: IMF (Industrial manufacture); PREP (Preparation)  
(polyoxometallate supported mixed metal oxide catalysts for conversion of alkanes)

RN 107-02-8 HCAPLUS  
CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
CN 2-Propenenitrile (9CI) (CA INDEX NAME)



RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 9 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
AN 2000:314395 HCAPLUS  
DN 132:308827  
TI Method for the removal of aldehydes from chemical-manufacturing product streams during distillative purification by the addition of electron-donating group-substituted aromatic amines prior to product

distillation  
 IN Patel, Natsu R.; Lewis, Vincent E.; Enderson, Margaret D.  
 PA Nalco/Exxon Energy Chemicals L.P., USA  
 SO Eur. Pat. Appl., 5 pp.  
 CODEN: EPXXDW

DT Patent  
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 999207	A2	20000510	EP 1999-307810	19991004
	EP 999207	A3	20000524		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	US 6074532	A	20000613	US 1998-186579	19981105
	JP 2000143606	A2	20000526	JP 1999-268036	19990922
	KR 2000035214	A	20000626	KR 1999-48479	19991104
PRAI	US 1998-186579	A	19981105		

AB A method is described for increasing the purification efficiency when distilling

off aldehyde contaminants during chemical-manufacturing processes by adding a substituted aromatic amine, having electron-donating substituents, and a catalytic quantity of a mineral acid, at the input of a distillation column where the impure product-containing stream is added to the distillation

column. The method is particularly useful for removal of aldehydes (e.g., acrolein) generated as a byproduct of acrylonitrile manufacture because the aldehyde forms a Schiff base with the aromatic amine which facilitates distillation of

the desired product (e.g., acrylonitrile). Suitable aromatic amines are 2-aminoaniline, 3,4-di-methylaniline, 4-ethylaniline, etc.

IC ICM C07C253-34

ICS C07C255-08

CC 35-2 (Chemistry of Synthetic High Polymers)

Section cross-reference(s): 23, 48

IT 107-13-1P, Acrylonitrile, preparation

RL: PNU (Preparation, unclassified); **PUR (Purification or recovery); PREP (Preparation)**

(method for the removal of aldehydes from chemical-manufacturing product streams

during distillative purification by the addition of electron-donating group-substituted aromatic amines prior to product distillation)

IT 75-07-0, Acetaldehyde, reactions 78-85-3, Methacrolein 100-52-7, Benzaldehyde, reactions 107-02-8, Acrolein, reactions 123-38-6, Propionaldehyde, reactions

RL: RCT (Reactant); **REM (Removal or disposal); PROC (Process); RACT (Reactant or reagent)**

(method for the removal of aldehydes from chemical-manufacturing product streams

during distillative purification by the addition of electron-donating group-substituted aromatic amines prior to product distillation)

IT 107-13-1P, Acrylonitrile, preparation

RL: PNU (Preparation, unclassified); **PUR (Purification or recovery); PREP (Preparation)**

(method for the removal of aldehydes from chemical-manufacturing product streams


during distillative purification by the addition of electron-donating group-substituted aromatic amines prior to product distillation)

RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



IT 107-02-8, Acrolein, reactions  
 RL: RCT (Reactant); REM (Removal or disposal); PROC (Process);  
 RACT (Reactant or reagent)  
 (method for the removal of aldehydes from chemical-manufacturing product streams during distillative purification by the addition of electron-donating group-substituted aromatic amines prior to product distillation)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 10 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2000:59123 HCAPLUS

DN 132:123030

TI Suppression of staining in manufacturing (meth)acrylonitriles

IN Kashihara, Kunio; Nakajima, Sadao

PA Hakuto K. K., Japan

SO Jpn. Kokai Tokkyo Koho, 4 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2000026391	A2	20000125	JP 1998-192814	19980708
PRAI	JP 1998-192814		19980708		

AB In the acrylonitrile process liquid which includes the acrolein and HCN, the nitroxyl radicals and acetic acid are used to suppress the staining. The acrylonitriles are manufactured by ammoxidn. of propylene. Thus, a mixture containing 200 g acrylonitrile, 0.1% HCN, and 2% acrolein was treated with 1.0 ppm 4-hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl and 0.5% AcOH to give a reaction mixture containing 12 ppm polymers.

IC ICM C07C253-32

ICS C07C253-26; C07C255-08

CC 35-2 (Chemistry of Synthetic High Polymers)

Section cross-reference(s): 23, 45

IT 64-19-7, Acetic acid, uses 2226-96-2,  
 4-Hydroxy-2,2,6,6-tetramethylpiperidine-1-oxyl 2564-83-2,  
 2,2,6,6-Tetramethylpiperidine-1-oxyl 2896-70-0, 4-Oxo-2,2,6,6-  
 tetramethylpiperidine-1-oxyl 14691-88-4 95407-69-5,  
 4-Methoxy-2,2,6,6-tetramethylpiperidine-1-oxyl

RL: CAT (Catalyst use); USES (Uses)

(polymerization inhibitor; suppression of staining in manufacturing (meth)acrylonitriles)

IT 74-90-8P, Hydrogen cyanide, preparation 107-02-8P, Acrolein, preparation

RL: BYP (Byproduct); PREP (Preparation)


(suppression of staining in manufacturing (meth)acrylonitriles)

IT 107-13-1P, Acrylonitrile, preparation


RL: IMF (Industrial manufacture); PREP (Preparation)

(suppression of staining in manufacturing (meth)acrylonitriles)

IT 64-19-7, Acetic acid, uses  
 RL: CAT (Catalyst use); USES (Uses)  
 (polymerization inhibitor; suppression of staining in manufacturing  
 (meth)acrylonitriles)  
 RN 64-19-7 HCAPLUS  
 CN Acetic acid (7CI, 8CI, 9CI) (CA INDEX NAME)



IT 107-02-8P, Acrolein, preparation  
 RL: BYP (Byproduct); PREP (Preparation)  
 (suppression of staining in manufacturing (meth)acrylonitriles)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



IT 107-13-1P, Acrylonitrile, preparation  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (suppression of staining in manufacturing (meth)acrylonitriles)  
 RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 11 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2000:43343 HCAPLUS  
 DN 132:93035  
 TI Method for purification of nitrile by removing aldehyde using anion  
 exchange resin  
 IN Takahashi, Shuya; Asai, Koich; Matsushita, Mitsuo; Uehara, Yoshikazu  
 PA Mitsui Chemicals Inc., Japan  
 SO Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2000016978	A2	20000118	JP 1998-184958	19980630
PRAI	JP 1998-184958		19980630		

AB Nitriles are purified by contacting the nitriles with anion exchange  
 resin-supported compound bearing active methylene group and acidic group in  
 the same mol. The above compound is preferably  $\alpha$ -substituted acetic  
 acid such as malonic acid, malonic acid monomethyl or monoethyl ester,  
 cyanoacetic acid, acetoacetic acid, sulfoacetic acid, or  
 acetonedicarboxylic acid. This process allows aldehydes to react with the  
 compound possessing active methylene group and acidic group and efficiently  
 and inexpensively removes minute quantity of aldehydes in nitriles. The  
 anion exchange resins used can be readily regenerated by the usual manner.  
 Thus, acrylonitrile containing 3 ppm acrolein was passed through a column of

15 mL Lewatit MP62 (Na+) malonic acid salt at 45 mL/h for 3 days to give acrylonitrile containing  $\leq 0.1$  ppm.

IC ICM C07C253-34  
ICS C07C255-08

CC 23-19 (Aliphatic Compounds)

IT 75-05-8P, Acetonitrile, preparation 107-13-1P, Acrylonitrile, preparation 126-98-7P, Methacrylonitrile  
RL: PUR (Purification or recovery); PREP (Preparation)  
(method for purification of nitrile by removing aldehyde using active hydrogen-containing acidic compound-carrying anion exchange resin)

IT 75-07-0, Acetaldehyde, processes 78-85-3, Methacrolein 107-02-8, Acrolein, processes  
RL: REM (Removal or disposal); PROC (Process)  
(method for purification of nitrile by removing aldehyde using active hydrogen-containing acidic compound-carrying anion exchange resin)

IT 107-13-1P, Acrylonitrile, preparation  
RL: PUR (Purification or recovery); PREP (Preparation)  
(method for purification of nitrile by removing aldehyde using active hydrogen-containing acidic compound-carrying anion exchange resin)

RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



IT 107-02-8, Acrolein, processes  
RL: REM (Removal or disposal); PROC (Process)  
(method for purification of nitrile by removing aldehyde using active hydrogen-containing acidic compound-carrying anion exchange resin)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 12 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 2000:43147 HCAPLUS

DN 132:79583

TI Polymers having active methylene groups for removal of aldehyde impurities in nitriles

IN Takahashi, Hidenari; Asai, Koichi; Matsushita, Mitsuo; Uehara, Yoshikazu

PA Mitsui Chemicals Inc., Japan

SO Jpn. Kokai Tokkyo Koho, 6 pp.  
CODEN: JKXXAF

DT Patent

LA Japanese


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2000015113	A2	20000118	JP 1998-182915	19980629
PRAI	JP 1998-182915		19980629		
OS	MARPAT 132:79583				
AB	Title polymers comprise anion-exchange polymers containing compds. having active methylene groups and acidic groups. Thus, a malonic acid-supported Lewatit MP 62 (weakly basic anion exchanger) removed acrolein in acrylonitrile effectively.				
IC	ICM B01J041-12				

ICS C08K005-09; C08K005-41; C08K005-49; C08L101-02; C07C253-34  
 CC 38-3 (Plastics Fabrication and Uses)  
 IT 75-05-8P, Acetonitrile, preparation 107-13-1P, Acrylonitrile,  
 preparation 126-98-7P, Methacrylonitrile  
 RL: PUR (Purification or recovery); PREP (Preparation)  
 (anion-exchange polymers containing compds. having active methylene groups  
 for removal of aldehydes in nitriles)  
 IT 75-07-0, Acetaldehyde, processes 78-85-3, Methacrolein 107-02-8  
 , Acrolein, processes  
 RL: REM (Removal or disposal); PROC (Process)  
 (anion-exchange polymers containing compds. having active methylene groups  
 for removal of aldehydes in nitriles)  
 IT 107-13-1P, Acrylonitrile, preparation  
 RL: PUR (Purification or recovery); PREP (Preparation)  
 (anion-exchange polymers containing compds. having active methylene groups  
 for removal of aldehydes in nitriles)  
 RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)




IT 107-02-8, Acrolein, processes  
 RL: REM (Removal or disposal); PROC (Process)  
 (anion-exchange polymers containing compds. having active methylene groups  
 for removal of aldehydes in nitriles)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 13 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1999:332838 HCAPLUS  
 DN 131:75241  
 TI Heterogeneous - catalytic gas - phase oxidation of  $\beta$ -picoline  
 AU Heinz, Dieter  
 CS Leverkusen, Germany  
 SO Fortschritt-Berichte VDI, Reihe 3: Verfahrenstechnik (1999), 582, I-VIII,  
 1-160  
 CODEN: FVVEFK; ISSN: 0178-9503  
 PB VDI Verlag GmbH  
 DT Journal  
 LA German  
 AB Expts. on the oxidation of  $\beta$ -picoline to nicotinic acid in the gas phase  
 using V2O5/TiO2 catalyst and the preparation of 3-cyanopyridine from acrolein,  
 acrylonitrile, and ammonia were described.  
 CC 45-4 (Industrial Organic Chemicals, Leather, Fats, and Waxes)  
 Section cross-reference(s): 67  
 IT 1314-23-4, Zirconium dioxide, uses 1314-62-1, Vanadium pentoxide, uses  
 13463-67-7, Titanium dioxide, uses  
 RL: CAT (Catalyst use); USES (Uses)  
 (catalyst; in gas-phase oxidation of  $\beta$ -picoline to nicotinic  
 acid)  
 IT 100-54-9, Nicotinonitrile 107-02-8, Acrolein, reactions  
 107-13-1, Acrylonitrile, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of nicotinonitrile from acrolein, acrylonitrile and ammonia)  
 IT 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile,  
 reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (preparation of nicotinonitrile from acrolein, acrylonitrile and ammonia)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 14 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1998:577249 HCAPLUS  
 DN 129:232294  
 TI Theoretical study of the alternative structure of copolymer ligands and  
 activity comparison of copolymer ligands  
 AU Li, Lei; Wang, Zuo-Xin; Wang, Xiao-Jun; Yuan, Guo-Qing  
 CS Beijing Univ. Chem. Technol., Beijing, 100029, Peop. Rep. China  
 SO Huaxue Xuebao (1998), 56(8), 747-754  
 CODEN: HHHPA4; ISSN: 0567-7351  
 PB Kexue Chubanshe  
 DT Journal  
 LA Chinese  
 AB A theor. study of the alternative structures of copolymer ligands (for Rh)  
 of catalyst for carbonylation of methanol to acetic acid has been carried  
 out in terms of the ASED-MO method and Monte Carlo method. AA, AB, BB, BA  
 reaction pathways were calculated and energy barriers were determined Based  
 on the  
 supposition of the preexponent factor of rate consts. being equal, the  
 reactivity ratios r1 and r2 were obtained. The copolymer structure was  
 simulated and the ratio of active AB segment in the copolymer mol. chains  
 was calculated The relative activity of copolymer ligand consisted of  
 different two units was compared. The effect of temperature change and the  
 feed  
 ratio x, on the ratio of AB in the mol. chains was also discussed.  
 CC 45-4 (Industrial Organic Chemicals, Leather, Fats, and Waxes)  
 Section cross-reference(s): 35, 67  
 IT 78-94-4, 3-Buten-2-one, reactions 79-10-7, Acrylic acid, reactions  
 96-33-3, Methyl acrylate 100-43-6, 4-Vinylpyridine 100-69-6,  
 2-Vinylpyridine 107-02-8, Acrolein, reactions 107-13-1  
 , Acrylonitrile, reactions 108-05-4, Vinyl acetate, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reactivity ratio in polymerization)  
 IT 7440-16-6D, Rhodium, complexes with vinylpyridine copolymers, uses  
 RL: CAT (Catalyst use); PRP (Properties); USES (Uses)  
 (theor. study of alternative structure and catalytic activity of  
 copolymer ligands in Rh complexes as catalysts for carbonylation of  
 methanol to acetic acid)  
 IT 25750-28-1D, Methyl acrylate-4-vinylpyridine copolymer, complexes with  
 rhodium 25750-29-2D, Acrylic acid-4-vinylpyridine copolymer,  
 complexes with rhodium 25852-53-3D, Acrylic acid



-2-vinylpyridine copolymer, complexes with rhodium 26836-60-2D, Acrylonitrile-2-vinylpyridine copolymer, complexes with rhodium 26877-88-3D, Vinyl acetate-4-vinylpyridine copolymer, complexes with rhodium 28408-26-6D, Vinyl acetate-2-vinylpyridine copolymer, complexes with rhodium 29729-87-1D, Methyl acrylate-2-vinylpyridine copolymer, complexes with rhodium 32236-74-1D, Acrylonitrile-4-vinylpyridine copolymer, complexes with rhodium 60292-97-9D, Acrolein-4-vinylpyridine copolymer, complexes with rhodium 90994-54-0D, complexes with rhodium 105133-74-2D, Butenone-2-vinylpyridine copolymer, complexes with rhodium 162844-26-0D, Acrolein-2-vinylpyridine copolymer, complexes with rhodium  
 RL: CAT (Catalyst use); PRP (Properties); USES (Uses)


(theor. study of alternative structure of copolymer ligands and catalytic activity of Rh complexes as catalysts for carbonylation of methanol to acetic acid)

IT 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile, reactions

RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reactivity ratio in polymerization)


RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 15 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1997:738775 HCAPLUS

DN 128:61167

TI High-pressure kinetics of Lewis acid catalyzed cycloadditions and ene reactions. A convincing method for mechanistic delineation

AU Jenner, G.

CS Laboratoire de Piezochimie Organique, Synthese et Stereoreactivite (CNRS URA 466), Institut de Chimie, Strasbourg, 67008, Fr.

SO New Journal of Chemistry (1997), 21(10), 1085-1090

CODEN: NJCHE5; ISSN: 1144-0546

PB Gauthier-Villars

DT Journal

LA English

AB The mechanistic course of Lewis acid catalyzed cycloaddns. and ene reactions is determined by high-pressure kinetics. On the basis of activation vols., the concertedness of [ $\pi_4 + \pi_2$ ] cycloaddns. is not altered by the presence of the catalyst. This is also valid for the [ $\pi_2 + \pi_2 + \pi_2$ ] cycloaddn., but not for the  $\text{AlCl}_3$  catalyzed cyclobutene formation in [2 + 2] cycloaddns., which proceeds stepwise. The Lewis acid catalyzed ene reaction involving (C-H-C) hydrogen transfer is a two-step process with a dipolar acyclic transition state. These results should give confidence in the high-pressure method to delineate reaction mechanisms.

CC 22-4 (Physical Organic Chemistry)

IT Lewis acids

RL: CAT (Catalyst use); USES (Uses)


(high-pressure kinetics of Lewis acid catalyzed cycloaddns. and ene reaction)

IT 7446-70-0, Aluminum trichloride, uses 7646-85-7, Zinc dichloride, uses 7791-03-9, Lithium perchlorate 10026-11-6, Zirconium tetrachloride 17631-68-4 18323-96-1  
 RL: CAT (Catalyst use); USES (Uses)  
 (high-pressure kinetics of Lewis acid catalyzed cycloaddns. and ene reaction)

IT 78-79-5, reactions 78-94-4, Methyl vinyl ketone, reactions 107-02-8, Acrolein, reactions 107-13-1, 2-Propenenitrile, reactions 109-92-2, Ethyl vinyl ether 110-83-8, Cyclohexene, reactions 121-46-0, Bicyclo[2.2.1]hepta-2,5-diene 498-66-8, Bicyclo[2.2.1]hept-2-ene 534-22-5, 2-Methylfuran 922-67-8, Methyl propiolate 4170-30-3, Crotonaldehyde  
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent)  
 (high-pressure kinetics of Lewis acid catalyzed cycloaddns. and ene reaction)

IT 107-02-8, Acrolein, reactions 107-13-1, 2-Propenenitrile, reactions  
 RL: PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent)  
 (high-pressure kinetics of Lewis acid catalyzed cycloaddns. and ene reaction)

RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



RE.CNT 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 16 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1997:617963 HCAPLUS  
 DN 127:283826  
 TI Functionalized nanotubes  
 IN Fischer, Alan; Hoch, Robert; Moy, David; Lu, Ming; Martin, Mark; Niu, Chun Ming; Ogata, Naoya; Tennent, Howard; Dong, Liwen; Sun, Ji; Helms, Larry; Jameison, Fabian; Liang, Pam; Simpson, David  
 PA Hyperion Catalysis International, Inc., USA  
 SO PCT Int. Appl., 133 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 5

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9732571	A1	19970912	WO 1997-US3553	19970305
W: AM, AT, AU, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, YU				
RW: GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB,				

GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN,  
ML, MR, NE, SN, TD, TG

CA 2247820	AA	19970912	CA 1997-2247820	19970305
AU 9721979	A1	19970922	AU 1997-21979	19970305
AU 724277	B2	20000914		
EP 910340	A1	19990428	EP 1997-914892	19970305
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
CN 1217653	A	19990526	CN 1997-194402	19970305
BR 9707845	A	19990727	BR 1997-7845	19970305
JP 2002503204	T2	20020129	JP 1997-531955	19970305
IL 125987	A1	20030212	IL 1997-125987	19970305
RU 2200562	C2	20030320	RU 1998-116596	19970305
PRAI US 1996-37238	P	19960306		
US 1996-37238P	P	19960306		
WO 1997-US3553	W	19970305		

AB Graphitic nanotubes, which include tubular fullerenes (commonly called buckytubes) and fibrils, which are functionalized by chemical substitution or by adsorption of functional moieties are claimed. More specifically the invention relates to graphitic nanotubes which are uniformly or nonuniformly substituted with chemical moieties or upon which certain cyclic compds. are adsorbed and to complex structures comprised of such functionalized nanotubes linked to one another. The invention also relates to methods for introducing functional groups onto the surface of such nanotubes. The invention further relates to uses for functionalized nanotubes, which include enzyme immobilization for sample separation and immobilizing a biocatalyst capable of catalyzing a reaction on the functionalized nanotubes.

IC ICM A61K009-00  
ICS A01N025-00; C09C001-56; B32B005-16

CC 66-4 (Surface Chemistry and Colloids)  
Section cross-reference(s): 7

IT 56-87-1DP, L-Lysine, carbon fibril bonded, preparation 58-85-5DP, Biotin, surface reaction product with carbon fibrils 60-24-2DP, Monothioethylene glycol, surface reaction product with carbon nanotubes and fibrils 75-89-8DP, 2,2,2-Trifluoroethanol, surface reaction product with carbon nanotubes and fibrils 79-06-1DP, 2-Propenamide, surface reaction product with carbon nanotubes and fibrils, preparation 79-10-7DP, 2-Propenoic acid, surface reaction product with carbon nanotubes and fibrils, preparation 107-02-8DP, Propenal, surface reaction product with carbon nanotubes and fibrils 107-11-9DP, 3-Amino-1-propene, surface reaction product with carbon nanotubes and fibrils 107-13-1DP, 2-Propenenitrile, surface reaction product with carbon nanotubes and fibrils, preparation 107-18-6DP, 2-Propen-1-ol, surface reaction product with carbon nanotubes and fibrils, preparation 108-31-6DP, 2,5-Furandione, surface reaction product with carbon nanotubes and fibrils, preparation 109-72-8DP, Butyllithium, surface reaction product with carbon nanotubes and fibrils 110-16-7DP, 2-Butenedioic acid (Z)-, surface reaction product with carbon nanotubes and fibrils 111-86-4DP, 1-Octanamine, surface reaction product with carbon nanotubes and fibrils 124-30-1DP, 1-Octadecanamine, surface reaction product with carbon nanotubes and fibrils 151-50-8DP, Potassium cyanide, surface reaction product with carbon nanotubes and fibrils 530-62-1DP, N,N'-Carbonyl diimidazole, surface reaction product with carbon nanotubes and fibrils 593-56-6DP, Methoxyamine hydrochloride, surface reaction product with carbon nanotubes and fibrils 814-68-6DP, Propenoyl chloride, surface reaction product with carbon nanotubes and fibrils 994-30-9DP, Chlorotriethylsilane, surface reaction product with carbon nanotubes and fibrils 1310-73-2DP, Sodium hydroxide, surface reaction product with carbon nanotubes and fibrils 1333-74-0DP,

Hydrogen, surface reaction product with carbon nanotubes and fibrils, preparation 1336-21-6DP, Ammonium hydroxide, surface reaction product with carbon nanotubes and fibrils 1892-57-5DP, 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide, surface reaction product with carbon nanotubes and fibrils 2016-57-1DP, 1-Aminodecane, surface reaction product with carbon nanotubes and fibrils 2074-87-5DP, Cyanogen, surface reaction product with carbon nanotubes and fibrils 4048-33-3DP, 6-Aminohexan-1-ol, surface reaction product with carbon nanotubes and fibrils 4781-83-3DP, 2-Iminothiolane hydrochloride, surface reaction product with carbon nanotubes and fibrils 5591-94-6DP, surface reaction product with carbon nanotubes and fibrils 5957-17-5DP, Triethyl(2-hydroxyethyl)ammonium iodide, surface reaction product with carbon nanotubes and fibrils 7664-41-7DP, Ammonia, surface reaction product with carbon nanotubes and fibrils, preparation 7664-93-9DP, Sulfuric acid, surface reaction product with carbon nanotubes and fibrils, preparation 7697-37-2DP, Nitric acid, surface reaction product with carbon nanotubes and fibrils, preparation 7704-34-9DP, Sulfur, surface reaction product with carbon nanotubes and fibrils, preparation 7732-18-5DP, Water, surface reaction product with carbon nanotubes and fibrils, preparation 7782-44-7DP, Oxygen, surface reaction product with carbon nanotubes and fibrils, preparation 13214-66-9DP, 4-Phenylbutylamine, surface reaction product with carbon nanotubes and fibrils 19008-71-0DP, 8-Aminooctan-1-ol, surface reaction product with carbon nanotubes and fibrils 23160-46-5DP, 10-Aminodecan-1-ol, surface reaction product with carbon nanotubes and fibrils 103708-09-4DP, Sulfosuccinimidyl-4-(N-maleimidomethyl)cyclohexanecarboxylate, surface reaction product with carbon nanotubes and fibrils 142755-63-3DP, 18-Aminooctadecan-1-ol, surface reaction product with carbon nanotubes and fibrils


RL: SPN (Synthetic preparation); PREP (Preparation)  
(surface functionalization of carbon nanotubes and fibrils for enzyme immobilization)

IT 60-24-2DP, Monothioethylene glycol, surface reaction product with carbon nanotubes and fibrils 107-02-8DP, Propenal, surface reaction product with carbon nanotubes and fibrils 107-13-1DP, 2-Propenenitrile, surface reaction product with carbon nanotubes and fibrils, preparation

RL: SPN (Synthetic preparation); PREP (Preparation)  
(surface functionalization of carbon nanotubes and fibrils for enzyme immobilization)


RN 60-24-2 HCAPLUS

CN Ethanol, 2-mercapto- (8CI, 9CI) (CA INDEX NAME)



RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)




L33 ANSWER 17 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1997:544238 HCAPLUS  
 DN 127:191191  
 TI Regeneration of weak basic anionic exchange resin for purification of acrylonitrile  
 IN Uehara, Yoshikazu; Kanbara, Yoshihiko; Tsukahara, Yasushi; Nozaki, Shohei  
 PA Mitsui Toatsu Chemicals, Inc., Japan  
 SO Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 09206604	A2	19970812	JP 1996-18657	19960205
PRAI	JP 1996-18657		19960205		
AB	The title process consists of contacting the used primary and/or secondary amine-containing exchange resins (e.g., Diaion WA-20, Lewatit S-100) having adsorbed aldehydes (e.g., acrolein) with mineral acids (e.g., H <sub>2</sub> SO <sub>4</sub> , HCl).				
IC	ICM B01J049-00				
CC	35-2 (Chemistry of Synthetic High Polymers) Section cross-reference(s): 46				
IT	107-13-1P, Acrylonitrile, preparation RL: <b>PUR (Purification or recovery); PREP (Preparation)</b> (regeneration of weak basic anionic exchange resin for purification of acrylonitrile)				
IT	107-02-8, Acrolein, processes RL: <b>REM (Removal or disposal); PROC (Process)</b> (regeneration of weak basic anionic exchange resin for purification of acrylonitrile)				
IT	107-13-1P, Acrylonitrile, preparation RL: <b>PUR (Purification or recovery); PREP (Preparation)</b> (regeneration of weak basic anionic exchange resin for purification of acrylonitrile)				
RN	107-13-1 HCAPLUS				
CN	2-Propenenitrile (9CI) (CA INDEX NAME)				




IT 107-02-8, Acrolein, processes  
 RL: **REM (Removal or disposal); PROC (Process)**  
 (regeneration of weak basic anionic exchange resin for purification of acrylonitrile)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 18 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1996:737386 HCAPLUS  
 DN 126:237943  
 TI Molybdovanadophosphate (NPMoV)/hydroquinone/O<sub>2</sub> system as an efficient reoxidation system in palladium-catalyzed oxidation of alkenes

- AU Yokota, Takahiro; Fujibayashi, Shinya; Nishiyama, Yutaka; Sakaguchi, Satoshi; Ishii, Yasutaka  
 CS Department of Applied Chemistry, Faculty of Engineering, Kansai University, Suita Osaka, 564, Japan  
 SO Journal of Molecular Catalysis A: Chemical (1996), 114(1-3), 113-122  
 CODEN: JMCCF2; ISSN: 1381-1169  
 PB Elsevier  
 DT Journal  
 LA English  
 OS CASREACT 126:237943  
 AB The molybdovanadophosphate (NPMoV)-hydroquinone (I)-O<sub>2</sub> system was an efficient reoxidn. system in Pd-catalyzed oxidns. of alkenes and related compds. Thus, acetoxylation of cycloalkenes with O<sub>2</sub> as the final oxidant were cleanly performed using the multicatalytic system consisting of Pd(OAc)<sub>2</sub>-I-NPMoV to form 3-acetoxy-1-cycloalkenes in good yields. Cyclopentene and cyclohexene were converted into the corresponding allylic acetates in almost quant. yields. Omitting I from the catalytic system led to low yields of the acetates. Acetoxylation of cyclooctene was achieved satisfactorily by replacing I in the multicatalytic system by chlorohydroquinone. Molybdovanadophosphates, which catalyze the smooth dehydrogenation of I to benzoquinone with O<sub>2</sub>, rapidly promoted the present Pd(II)-catalyzed acetoxylation of cycloalkenes. By the use of a mixed EtOH-H<sub>2</sub>O solvent under these conditions, Wacker-type oxidns. of cyclohexene and styrene were accomplished in fair-to-good yields. Monosubstituted alkenes such as Et acrylate and acrylonitrile underwent acetalization by this catalytic system to give the corresponding acetals quant.
- CC 21-2 (General Organic Chemistry)  
 Section cross-reference(s): 67
- IT 123-31-9, Hydroquinone, uses 3375-31-3, Palladium diacetate 7631-95-0, Disodium molybdate 7664-38-2, Phosphoric acid, uses 7782-44-7, Oxygen, uses 13718-26-8, Sodium metavanadate  
 RL: CAT (Catalyst use); USES (Uses)  
 (molybdovanadophosphate (NPMoV)-hydroquinone-O<sub>2</sub> system as efficient reoxidn. system in palladium-catalyzed oxidation of alkenes)
- IT 100-42-5, Styrene, reactions 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile, reactions 110-83-8, Cyclohexene, reactions 140-88-5, Ethyl acrylate 142-29-0, Cyclopentene 591-47-9, 4-Methyl-1-cyclohexene 591-49-1 628-92-2, Cycloheptene 931-88-4, Cyclooctene 1501-82-2, Cyclododecene 3054-95-3, Acrolein diethyl acetal 38451-18-2  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (molybdovanadophosphate (NPMoV)-hydroquinone-O<sub>2</sub> system as efficient reoxidn. system in palladium-catalyzed oxidation of alkenes)
- IT 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (molybdovanadophosphate (NPMoV)-hydroquinone-O<sub>2</sub> system as efficient reoxidn. system in palladium-catalyzed oxidation of alkenes)
- RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



- RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



RE.CNT 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L33 ANSWER 19 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1995:574914 HCAPLUS

DN 123:40057

TI The development of a novel strategy for the microbial treatment of acrylonitrile effluents

AU Wyatt, J.M.; Knowles, C.J.

CS Biological Laboratory, University of Kent, Canterbury, CT2 7NJ, UK

SO Biodegradation (1995), 6(2), 93-107

CODEN: BIODG; ISSN: 0923-9820

DT Journal

LA English

AB Effluent from acrylonitrile manufacturing is difficult to biodegrade. It contains 9 major organic components: acetic acid, acrylonitrile, acrylamide, acrylic acid, acrolein, cyanopyridine, fumaronitrile, succinonitrile, and maleimide. A range of bacteria were isolated that can grow on, or convert all the organic components of acrylonitrile manufacturing effluent. These bacteria

can be used as the basis of a mixed culture system to treat the effluent. The bacteria were utilized in batch and continuous cultures to degrade a synthetic wastewater containing acrylonitrile, acrylamide, acrylic acid, cyanopyridine, and succinonitrile. The mixed microbial population was adapted by varying growth rates and switching from continuous to batch and back to continuous growth, to degrade these 5 compds. and acrolein, fumaronitrile, and maleimide.

CC 60-1 (Waste Treatment and Disposal)

Section cross-reference(s): 10

IT 107-13-1P, Acrylonitrile, processes

RL: BPR (Biological process); BSU (Biological study, unclassified); IMF (Industrial manufacture); REM (Removal or disposal); BIOL (Biological study); PREP (Preparation); PROC (Process)

(strategy for mixed culture bacterial degradation of organic components of acrylonitrile manufacturing effluent)

IT 79-06-1, Acrylamide, processes 79-10-7, Acrylic acid, processes

107-02-8, Acrolein, processes 110-61-2, Succinonitrile

541-59-3, Maleimide 764-42-1, Fumaronitrile 29386-66-1, Cyanopyridine

RL: BPR (Biological process); BSU (Biological study, unclassified);

REM (Removal or disposal); BIOL (Biological study); PROC (Process)

(strategy for mixed culture bacterial degradation of organic components of acrylonitrile manufacturing effluent)

IT 107-13-1P, Acrylonitrile, processes

RL: BPR (Biological process); BSU (Biological study, unclassified); IMF (Industrial manufacture); REM (Removal or disposal); BIOL (Biological study); PREP (Preparation); PROC (Process)

(strategy for mixed culture bacterial degradation of organic components of acrylonitrile manufacturing effluent)

RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)




IT 107-02-8, Acrolein, processes

RL: BPR (Biological process); BSU (Biological study, unclassified);

REM (Removal or disposal); BIOL (Biological study); PROC (Process)  
 (strategy for mixed culture bacterial degradation of organic components of  
 acrylonitrile manufacturing effluent)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 20 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1994:458200 HCAPLUS

DN 121:58200

TI Ammoxidation of olefins

IN Paparizos, Christos; Shaw, Wilfrid Garside

PA Standard Oil Co., USA

SO Eur. Pat. Appl., 10 pp.

CODEN: EPXXDW

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 573713	A1	19931215	EP 1992-305436	19920612
	EP 573713	B1	19970102		
	R: AT, DE, ES, GB, IT, NL				
	US 5134105	A	19920728	US 1990-495875	19900319
	AT 147070	E	19970115	AT 1992-305436	19920612
	ES 2095406	T3	19970216	ES 1992-305436	19920612
PRAI	US 1990-495875		19900319		
	EP 1992-305436	A	19920612		

AB Olefins such as propylene and isobutylene are converted to the corresponding unsatd. nitriles, acrylonitrile, and methacrylonitrile, resp., by reacting a mixture of the olefin, ammonia, and mol. oxygen-containing gas in the presence of a catalyst containing the oxides of molybdenum, bismuth, iron, cobalt, nickel, and chromium, and either phosphorus or antimony or mixts. thereof, and an alkali metal or mixture thereof, and optionally one element selected from the group of an alkaline earth metal, a rare earth metal, niobium, thallium, arsenic, magnesium, zinc, cadmium, vanadium, boron, calcium, tin, germanium, manganese, tungsten, tellurium, or mixts. thereof.

IC ICM C07C253-26

ICS C07C255-08


CC 35-2 (Chemistry of Synthetic High Polymers)

Section cross-reference(s): 67

IT 1309-64-4, Antimony trioxide, uses 1333-82-0, Chromium trioxide  
 7439-95-4D, Magnesium, complex oxides 7439-96-5D, Manganese, complex  
 oxides 7440-03-1D, Niobium, complex oxides 7440-28-0D, Thallium,  
 complex oxides 7440-31-5D, Tin, complex oxides 7440-33-7D, Tungsten,  
 complex oxides 7440-38-2D, Arsenic, complex oxides 7440-42-8D, Boron,  
 complex oxides 7440-43-9D, Cadmium, complex oxides 7440-56-4D,  
 Germanium, complex oxides 7440-62-2D, Vanadium, complex oxides  
 7440-66-6D, Zinc, complex oxides 7440-70-2D, Calcium, complex oxides  
 7664-38-2, Phosphoric acid, uses 7757-79-1, Nitric  
 acid potassium salt, uses 7782-61-8, Ferric nitrate nonahydrate  
 7789-18-6, Cesium nitrate 10026-22-9, Cobalt nitrate hexahydrate  
 10035-06-0, Bismuth nitrate pentahydrate 12027-67-7, Ammonium  
 heptamolybdate 13478-00-7, Nickel nitrate hexahydrate 13494-80-9D,  
 Tellurium, complex oxides 62067-70-3 155948-76-8 155948-78-0



156260-89-8 156260-90-1  
 RL: CAT (Catalyst use); USES (Uses)  
 (catalysts, for ammoxidn. of olefins to nitriles)  
 IT 107-02-8P, 2-Propenal, preparation 107-13-1P,  
 2-Propenenitrile, preparation  
 RL: PREP (Preparation)  
 (preparation of, by ammoxidn. of propylene, catalysts for)  
 IT 107-02-8P, 2-Propenal, preparation 107-13-1P,  
 2-Propenenitrile, preparation  
 RL: PREP (Preparation)  
 (preparation of, by ammoxidn. of propylene, catalysts for)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)




RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 21 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1992:650942 HCAPLUS  
 DN 117:250942  
 TI Homogeneous catalytic hydrogenation of olefins using  $\text{RhH}_2(\text{Ph}_2\text{N}_3)(\text{PPh}_3)_2$  in tetrahydrofuran  
 AU Kameda, Noriyuki; Igarashi, Reiko  
 CS Nihom Univ., Funabashi, 274, Japan  
 SO Journal of Molecular Catalysis (1992), 75(1), 15-20  
 CODEN: JMCADS; ISSN: 0304-5102  
 DT Journal  
 LA English  
 OS CASREACT 117:250942  
 AB The catalytic hydrogenation activity of  $\text{RhH}_2(\text{Ph}_2\text{N}_3)(\text{PPh}_3)_2$  (I) in THF with different unsatd. compds. (allyl alc., cinnamic alc., cinnamaldehyde, cinnamic acid, cinnamionitrile, acrylic acid, acrolein, acrylonitrile, and styrene) was studied. Acrolein, acrylic acid, and acrylonitrile were reduced in the presence of I, whereas, other substrates were not. The highest activity was observed with acrylonitrile. A possible mechanism for this reaction is suggested.  
 CC 23-19 (Aliphatic Compounds)  
 Section cross-reference(s): 22  
 IT 36059-83-3  
 RL: CAT (Catalyst use); USES (Uses)  
 (catalysts, for hydrogenation of acrylonitrile, acrolein, and acrylic acid)  
 IT 79-10-7, Acrylic acid, reactions 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (hydrogenation of, in presence of rhodium hydride complex)  
 IT 107-02-8, Acrolein, reactions 107-13-1, Acrylonitrile, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (hydrogenation of, in presence of rhodium hydride complex)  
 RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 22 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1991:229693 HCAPLUS

DN 114:229693

TI Process for the manufacture of vinylcarbonyl compound-modified polyoxyalkylenes and their derivatives as reactive intermediates

IN Kuehn, Manfred

PA Akademie der Wissenschaften der DDR, Zentralinstitut fuer Molekularbiologie, Ger. Dem. Rep.

SO Ger. (East), 4 pp.

CODEN: GEXXA8

DT Patent

LA German

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DD 284239	A5	19901107	DD 1989-328842	19890524
PRAI	DD 1989-328842		19890524		

AB The title compds., useful as active intermediates for biotechnol., biochem., chemical, pharmaceuticals, and in medicines (no data), are prepared by the reaction of polyoxyalkylenes and their monoalkoxy derivs. (as well as amino group- or mercapto group-containing derivs.) in the presence of a basic or acid catalyst in aqueous solution, organic solution, or their mixture at a

20-150° with a vinyl carbonyl compound  $\text{R}_1\text{CH}:\text{C}(\text{R}_2)\text{R}_3$  ( $\text{R}_1 = \text{H}, \text{Me}, \text{Ph}, \text{C}_6\text{H}_4\text{NO}_2$ ;  $\text{R}_2 = \text{H}, \text{methyl}$ ;  $\text{R}_3 = \text{CN}, \text{CHO}, \text{CO}_2\text{H}, \text{CONH}_2, \text{CONHNH}_2, \text{CO}_2\text{X}$ ;  $\text{X} = \text{C}_1\text{-4 alkyl}$ ). Thus, 3 g  $\omega, \omega'$ -diaminopolyethylene glycol (mol. weight 6000) was dissolved in the mixture of 40 mL chloroform and 10 mL ethanol, 1 mL acrylonitrile dissolved in 5 mL ethanol added, the mixture first stirred for 1 h at room temperature, distribution 8 h at 60°, 50 mL  $\text{H}_2\text{O}$  added, the organic phase separated, washed twice with equal amts. of  $\text{H}_2\text{O}$ , dried with  $\text{Na}_2\text{SO}_4$ , and the chloroform evaporated, producing 2.1 g of a nitrile group-modified polyethylene glycol derivative residue.

IC ICM C08G065-00

CC 35-8 (Chemistry of Synthetic High Polymers)

IT 64-19-7, Acetic acid, uses and miscellaneous 100-85-6, Benzyltrimethylammonium hydroxide 109-63-7, Boron trifluoride etherate 584-08-7, Potassium carbonate 7446-70-0, Aluminum chloride, uses and miscellaneous 7637-07-2, Boron trifluoride, uses and miscellaneous 7646-85-7, Zinc chloride ( $\text{ZnCl}_2$ ), uses and miscellaneous 7664-93-9, Sulfuric acid, uses and miscellaneous

RL: CAT (Catalyst use); USES (Uses)

(catalyst, for reaction of vinylcarbonyl compds. with polyoxyalkylenes and their derivs.)

IT 79-10-7DP, 2-Propenoic acid, reaction products with polyoxyalkylenes

80-62-6DP, reaction products with polyoxyalkylenes 104-55-2DP,

Cinnamaldehyde, reaction products with polyoxyalkylenes 107-02-8DP

, Acrolein, reaction products with polyoxyalkylenes 107-13-1DP,  
 2-Propenenitrile, reaction products with polyoxyalkylenes 4170-30-3DP,  
 Crotonaldehyde, reaction products with polyoxyalkylenes 9004-74-4DP,  
 Polyethylene glycol monomethylether, reaction products with vinylcarbonyl  
 compds. 24991-53-5DP, reaction products with vinylcarbonyl compds.  
 25322-68-3DP, reaction products with vinylcarbonyl compds. 68865-60-1DP,  
 reaction products with vinylcarbonyl compds. 80506-64-5DP, reaction  
 products with vinylcarbonyl compds.

RL: IMF (Industrial manufacture); PREP (Preparation)

(manufacture of, as reactive intermediates for biotechnol. and biochem. and  
 chemical and pharmaceuticals and medicine, catalyst for)


IT 64-19-7, Acetic acid, uses and miscellaneous

RL: CAT (Catalyst use); USES (Uses)

(catalyst, for reaction of vinylcarbonyl compds. with polyoxyalkylenes  
 and their derivs.)

RN 64-19-7 HCAPLUS

CN Acetic acid (7CI, 8CI, 9CI) (CA INDEX NAME)




IT 107-02-8DP, Acrolein, reaction products with polyoxyalkylenes  
 107-13-1DP, 2-Propenenitrile, reaction products with  
 polyoxyalkylenes

RL: IMF (Industrial manufacture); PREP (Preparation)

(manufacture of, as reactive intermediates for biotechnol. and biochem. and  
 chemical and pharmaceuticals and medicine, catalyst for)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 23 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1985:148822 HCAPLUS

DN 102:148822

TI Reductive arylation of electron-deficient olefins: 4-(4-  
 chlorophenyl)butan-2-one (2-butanone, 4-(4-chlorophenyl)-)

AU Citterio, Attilio

CS Ist. Chim., Politec. Milano, Milan, I-20133, Italy


SO Organic Syntheses (1984), 62, 67-73

CODEN: ORSYAT; ISSN: 0078-6209


DT Journal

LA English

GI



- AB Benzenediazonium compds. 4-RC<sub>6</sub>H<sub>4</sub>N<sub>2</sub><sup>+</sup> (R = H, Br, Cl, OMe, MeCO) were treated with R<sub>1</sub>CR<sub>2</sub>:CHR<sub>3</sub> (R<sub>1</sub> = H, Me, CHMe<sub>2</sub>, CMe<sub>3</sub>, Ph; R<sub>2</sub> = H, Me; R<sub>3</sub> = COMe, CHO, cyano, CO<sub>2</sub>H, CO<sub>2</sub>Et) and TiCl<sub>3</sub> to yield reductive arylation products I. Thus, CH<sub>2</sub>:CHCOME was added to TiCl<sub>3</sub> in DMF under N, 4-ClC<sub>6</sub>H<sub>4</sub>N<sub>2</sub><sup>+</sup> Cl<sup>-</sup> was introduced slowly, and the mixture was stirred to give 4-ClC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>CH<sub>2</sub>COME.
- CC 25-16 (Benzene, Its Derivatives, and Condensed Benzenoid Compounds)  
Section cross-reference(s): 23
- IT 7705-07-9, uses and miscellaneous  
RL: CAT (Catalyst use); USES (Uses)  
(catalyst, for reductive arylation of vinyl ketones, acrolein and acrylic acid by benzenediazonium salts)
- IT 79-10-7, reactions 107-02-8, reactions 107-13-1,  
reactions 122-57-6 140-88-5 141-79-7 625-33-2 5166-53-0  
26465-92-9  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reductive arylation of, by chlorobenzenediazonium chloride, catalyst for)
- IT 107-02-8, reactions 107-13-1, reactions  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(reductive arylation of, by chlorobenzenediazonium chloride, catalyst for)
- RN 107-02-8 HCAPLUS
- CN 2-Propenal (9CI) (CA INDEX NAME)



- RN 107-13-1 HCAPLUS
- CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 24 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1984:491632 HCAPLUS

DN 101:91632

TI Removal of acrolein from acrylonitrile product streams

IN Denicola, Thomas Kevin

PA Monsanto Co. , USA

SO Eur. Pat. Appl., 11 pp.  
CODEN: EPXXDW

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 110861	A1	19840613	EP 1983-870122	19831118
	EP 110861	B1	19860730		
	R: DE, FR, GB, IT, NL				

PRAI US 1982-443356 A 19821122

AB Acrolein (I) [107-02-8] is removed from a crude acrylonitrile (II) [107-13-1] product stream in a recovery column by maintaining the pH at 5.25-7 in a zone of maximum I concentration of the column. Thus, a stream of

II containing 229 ppm acrolein in a recovery column (column bottom temperature 73°, column top temperature 83°, pressure 13.8 kPa) was injected with NaOH at tray 18 of 40 trays to adjust pH to 6.5, and the I content in the stream was reduced to 7 ppm.

IC C07C121-32

CC 35-2 (Chemistry of Synthetic High Polymers)

IT 107-13-1P, uses and miscellaneous

RL: PREP (Preparation); USES (Uses)

(acrolein removal from, pH effect on)

IT 107-02-8, uses and miscellaneous

RL: REM (Removal or disposal); PROC (Process)

(removal of, from acrylonitrile, pH effect on)

IT 107-13-1P, uses and miscellaneous

RL: PREP (Preparation); USES (Uses)

(acrolein removal from, pH effect on)

RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)




IT 107-02-8, uses and miscellaneous

RL: REM (Removal or disposal); PROC (Process)

(removal of, from acrylonitrile, pH effect on)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 25 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1983:613024 HCAPLUS

DN 99:213024

TI Removing aldehyde impurities from acrylonitrile and acrylamide

IN Schmitt, Joseph Michael

PA American Cyanamid Co. , USA

SO Ger. Offen., 12 pp.

CODEN: GWXXBX

DT Patent

LA German

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 3241198	A1	19830804	DE 1982-3241198	19821108
	GB 2114118	A1	19830817	GB 1982-28822	19821008
	FR 2520738	A1	19830805	FR 1982-21342	19821220
	JP 58134063	A2	19830810	JP 1983-9483	19830125
	NL 8300333	A	19830816	NL 1983-333	19830128
PRAI	US 1982-344087	A	19820129		

AB Aldehyde impurities are removed from acrylamide [79-06-1] and acrylonitrile (I) [107-13-1] using a weakly basic ion exchange resin containing primary or secondary amines. Thus, I containing approx. 20-25 ppm

acrolein (II) [107-02-8] was passed through Amberlite IRA-45 [9055-98-5] ion-exchange resin at 10, 30, and 300 mL/h (contact time 30, 10, or 1 min, resp.) to give I containing <1 ppm II.

IC C07C121-32; C07C103-133; C07C120-00; C07C102-00  
 CC 35-2 (Chemistry of Synthetic High Polymers)  
 IT 79-06-1P, preparation 107-13-1P, preparation  
 RL: PREP (Preparation)  
 (acrolein removal from, with ion-exchange resin)

IT 107-02-8, uses and miscellaneous  
 RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylamide and acrylonitrile, ion-exchange resins in)


IT 107-13-1P, preparation  
 RL: PREP (Preparation)  
 (acrolein removal from, with ion-exchange resin)

RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



IT 107-02-8, uses and miscellaneous  
 RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylamide and acrylonitrile, ion-exchange resins in)

RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 26 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1982:527083 HCAPLUS

DN 97:127083

TI Hydrocyanation of activated olefins

IN Pesa, Frederick Anthony; Graham, Anne Marie

PA Standard Oil Co., USA

SO Eur. Pat. Appl., 21 pp.

CODEN: EPXXDW

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 51697	A1	19820519	EP 1980-303540	19801008
	EP 51697	B1	19840208		
	R: BE, CH, DE, FR, GB, IT, LU, NL, SE				
	CA 1151669	A1	19830809	CA 1980-361379	19801002
PRAI	EP 1980-303540		19801008		

OS CASREACT 97:127083


AB The reaction of RCR1:CR2R3 (R, R1, and R2 independently are H, alkyl, aryl, cycloalkyl, aralkyl, CO2H, carbalkoxy, alkanoyl, carboxyalkyl, carbalkoxyalkyl, alkanoylalkyl; R3 = CO2H, carbalkoxy, cyano, alkanoyl, carboxyalkyl, carbalkoxyalkyl, cyanoalkyl, alkanoylalkyl) with HCN was catalyzed by group IA and IIA elements. The hydrocyanation of CH2:CHCO2Me over LiOAc gave 80.4% MeO2CCH2CH2CN.

IC C07C120-02; C07C121-407; C07C121-34; C07C121-20; B01J023-02; B01J023-04

CC 23-19 (Aliphatic Compounds)

IT 7757-79-1, uses and miscellaneous

RL: CAT (Catalyst use); USES (Uses)  
 (catalysts, for hydrocyanation of Me acrylate and acrylic acid  
 )  
 IT 79-10-7, reactions 96-33-3 107-02-8, reactions  
 107-13-1, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (hydrocyanation of, catalysts for)  
 IT 107-02-8, reactions 107-13-1, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (hydrocyanation of, catalysts for)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 27 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1980:494826 HCAPLUS

DN 93:94826

TI Purification of acrylonitrile

IN Saito, Jun; Hara, Junji; Mitsuishi, Takatoshi; Fujii, Kenzo

PA Mitsui Toatsu Chemicals, Inc., Japan

SO Jpn. Kokai Tokkyo Koho, 5 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 55013205	A2	19800130	JP 1978-78523	19780630
	JP 60051458	B4	19851114		
PRAI	JP 1978-78523	A	19780630		

AB H<sub>2</sub>C:CHCN (I) was purified by contacting the liquid phase of I with aqueous phases (pH 7.0-9.0) containing amine salts. Thus, 10 mL com. I containing 2.5 ppm

H<sub>2</sub>C:CHCHO (II) was shaken with 10 mL of an aqueous amine solution (prepared by adding 10 g EtNH<sub>2</sub>.HCl to 80 g H<sub>2</sub>O, adjusting to pH 8.0 with aqueous NaOH, and diluting to 100 mL with H<sub>2</sub>O) 3 h to leave as low as <0.1 ppm II and <5 ppm EtNHCH<sub>2</sub>CH<sub>2</sub>CN in I.

IC C07C121-32; C07C120-00

CC 23-19 (Aliphatic Compounds)

IT 107-13-1P, preparation

RL: PUR (Purification or recovery); PREP (Preparation)  
 (purification of)

IT 107-02-8, uses and miscellaneous

RL: REM (Removal or disposal); PROC (Process)  
 (removal of, in purification of acrylonitrile)

IT 107-13-1P, preparation


RL: PUR (Purification or recovery); PREP (Preparation)  
 (purification of)

RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



IT 107-02-8, uses and miscellaneous  
 RL: REM (Removal or disposal); PROC (Process)  
 (removal of, in purification of acrylonitrile)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 28 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1980:129606 HCAPLUS  
 DN 92:129606  
 TI Purification of acrylonitrile  
 IN Saito, Jun; Hara, Junji; Mitsuishi, Takatoshi; Fujii, Kenzo  
 PA Mitsui Toatsu Chemicals, Inc., Japan  
 SO Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 54151915	A2	19791129	JP 1978-60040	19780522
	JP 58001108	B4	19830110		
PRAI	JP 1978-60040	A	19780522		

AB Acrylonitrile (I) [107-13-1] was purified by contacting liquid I with porous ion exchangers containing primary and secondary amino exchangeable groups. Thus, com. I containing 2.5 ppm acrolein (II) [107-02-8] was treated with 50 mL porous Diaion WA 20 [12619-71-5] column (1.7 + 22 cm) at 200 mL/h for 5 h, thereby reducing the II content to <0.1 ppm.

IC C07C121-32

CC 35-2 (Synthetic High Polymers)

IT 107-13-1P, preparation

RL: PUR (Purification or recovery); PREP (Preparation)  
 (purification of, amino group-containing porous ion exchanger for)

IT 107-02-8, uses and miscellaneous

RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylonitrile, amino group-containing porous ion exchanger for)

IT 107-13-1P, preparation

RL: PUR (Purification or recovery); PREP (Preparation)  
 (purification of, amino group-containing porous ion exchanger for)

RN 107-13-1 HCAPLUS


CN 2-Propenenitrile (9CI) (CA INDEX NAME)



IT 107-02-8, uses and miscellaneous  
 RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylonitrile, amino group-containing porous ion



exchanger for)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 29 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1976:74851 HCAPLUS  
 DN 84:74851  
 TI Purification of acrylonitrile  
 IN Frost, Brian W.  
 PA BP Chemicals International Ltd., UK  
 SO Fr. Demande, 5 pp.  
 CODEN: FRXXBL  
 DT Patent  
 LA French  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	FR 2256914	A1	19750801	FR 1975-199	19750106
	GB 1431511	A	19760407	GB 1974-668	19741219
	CA 1019346	A1	19771018	CA 1974-217057	19741230
PRAI	GB 1974-668	A	19740107		

AB Acrolein (I) [107-02-8] and HCN [74-90-8] impurities in acrylonitrile (II) [107-13-1], prepared by catalytic reaction of I with O and NH<sub>3</sub> at high temperature, were removed by treating impure II with an aqueous solution of an alkali metal, alkaline earth metal, or ammonium salt of an aliphatic carboxylic acid. Thus, 100 ml of a 1500 g/l. aqueous NH<sub>4</sub>OAc [631-61-8] solution was mixed at 38° with 100 ml of a II solution containing 5000 ppm I and 100 ppm succinonitrile (III). After 200 min the I concentration was reduced to 5 ppm and the III concentration was unchanged, confirming that no secondary reactions occurred.

IC C07C  
 CC 35-2 (Synthetic High Polymers)  
 IT 107-13-1P, preparation  
 RL: PUR (Purification or recovery); PREP (Preparation)  
 (purification of, ammonium acetate treatment for removal of acrolein and hydrocyanic acid in)

IT 107-02-8, uses and miscellaneous  
 RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylonitrile, ammonium acetate treatment for)


IT 107-13-1P, preparation  
 RL: PUR (Purification or recovery); PREP (Preparation)  
 (purification of, ammonium acetate treatment for removal of acrolein and hydrocyanic acid in)

RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)




IT 107-02-8, uses and miscellaneous  
 RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylonitrile, ammonium acetate treatment for)

RN 107-02-8 HCAPLUS  
CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 30 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
AN 1973:518720 HCAPLUS  
DN 79:118720  
TI Gas-chromatography studies of physicochemical properties of  
bismuth-molybdenum catalysts  
AU Lewicki, Andrzej; Paryjczak, Tadeusz; Beres, Janusz  
CS Politech. Lodz, Lodz, Pol.  
SO Roczniki Chemii (1973), 47(5), 981-7  
CODEN: ROCHAC; ISSN: 0035-7677  
DT Journal  
LA Polish  
AB The sorption of MeCH:CH<sub>2</sub>, CH<sub>2</sub>:CHCHO, CH<sub>2</sub>:CHCO<sub>2</sub>H, AcH, AcOH, and CH<sub>2</sub>:CHCN  
on Bi<sub>2</sub>O<sub>3</sub>, MoO<sub>3</sub>, Bi<sub>2</sub>O<sub>3</sub>.3MoO<sub>3</sub> (α-phase), Bi<sub>2</sub>O<sub>3</sub>.2MoO<sub>3</sub> (β-phase),  
and Bi<sub>2</sub>O<sub>3</sub>.MoO<sub>3</sub> (γ-phase) was studied by gas chromatog. at  
50-350°. Except for CH<sub>2</sub>:CHCO<sub>2</sub>H, the retention volume of all the  
substances studied increased slightly with increasing temperature The highest  
sorpitive properties were observed for the γ and the α + γ  
phases.  
CC 67-1 (Catalysis and Reaction Kinetics)  
Section cross-reference(s): 66  
IT 1304-76-3 13565-96-3 13595-85-2 16229-40-6  
RL: CAT (Catalyst use); USES (Uses)  
(catalysts, sorption by, of acrylic acid and related polar  
organic mols.)  
IT 64-19-7, properties 75-07-0, properties 79-10-7, properties  
107-02-8, properties 107-13-1, properties 115-07-1,  
properties  
RL: PEP (Physical, engineering or chemical process); PROC (Process)  
(sorption of, by bismuth oxide-molybdenum oxide catalysts)  
IT 64-19-7, properties 75-07-0, properties 79-10-7, properties  
107-02-8, properties 107-13-1, properties 115-07-1,  
properties  
RL: PEP (Physical, engineering or chemical process); PROC (Process)  
(sorption of, on bismuth molybdenum oxide catalysts)  
IT 107-02-8, properties 107-13-1, properties  
RL: PEP (Physical, engineering or chemical process); PROC (Process)  
(sorption of, by bismuth oxide-molybdenum oxide catalysts)  
RN 107-02-8 HCAPLUS  
CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
CN 2-Propenenitrile (9CI) (CA INDEX NAME)




IT 107-02-8, properties 107-13-1, properties

RL: PEP (Physical, engineering or chemical process); PROC (Process)  
(sorption of, on bismuth molybdenum oxide catalysts)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 31 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1972:487917 HCAPLUS

DN 77:87917

TI Catalysts for the oxidation of propylene to acrylonitrile or acrolein

IN Fattore, Vittorio; Notari, Bruno

PA Snamprogetti SpA

SO Ger. Offen., 15 pp. Addn. to Ger. 2,117,351.

CODEN: GWXXBX

DT Patent

LA German

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 2165978	A	19720622	DE 1971-2165978	19710408
	NO 130042	B	19740701	NO 1971-1299	19710405
	DK 132605	B	19760112	DK 1971-1708	19710407
	FR 2093436	A5	19720128	FR 1971-12678	19710409
	BE 765621	A1	19710830	BE 1971-3011	19710413
	CH 530378	A	19721115	CH 1971-530378	19710413
	AT 313866	B	19740311	AT 1971-3114	19710413
	HU 165998	P	19741228	HU 1971-SA2187	19710413
	PL 83201	P	19751231	PL 1971-147457	19710413
	CA 981241	A1	19760106	CA 1971-110230	19710413
	CS 174813	P	19770429	CS 1971-2630	19710413
	NL 7104954	A	19711018	NL 1971-4954	19710414
	ZA 7102388	A	19720126	ZA 1971-2388	19710414
	ES 390536	A1	19740401	ES 1971-390536	19710414
	SU 425380	D	19740425	SU 1971-1641109	19710414
	RO 56406	P	19740901	RO 1971-66575	19710414
	US 3850975	A	19741126	US 1971-133886	19710414
	JP 51040557	B4	19761104	JP 1971-23209	19710414
	SE 399704	C	19780608	SE 1971-4859	19710414
	GB 1354152	A	19740522	GB 1971-26693	19710419
	GB 1354153	A	19740522	GB 1973-39096	19710419
	PL 88981	P	19761030	PL 1972-153372	19720208
	US 3983054	A	19760928	US 1974-492222	19740726
	SE 7609779	A	19760903	SE 1976-9779	19760903
PRAI	IT 1970-23270	A	19700414		
	US 1971-133886	A3	19710414		

AB Addition to Ger. 2,117,351. U oxide-Te oxide catalysts of 1:4 U-Te molar ratio, optionally containing 25-50% SiO<sub>2</sub> as support, were prepared by evaporating or

spray drying, then calcining the mixed aqueous solns. containing UO<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O

(I),  $\text{H}_2\text{TeO}_4 \cdot 2\text{H}_2\text{O}$  (II), and optionally  $\text{SiO}_2$  solution. The catalysts were used in a fluidized bed for the preparation of  $\text{CH}_2\text{:CHCN}$  (III) or  $\text{CH}_2\text{:CHCHO}$  by oxidation

of  $\text{C}_3\text{H}_6$  with air in the presence of  $\text{NH}_3$  or steam, resp., to give  $\text{C}_3\text{H}_6$  conversion 36-84.3 mole % and III selectivity 53.4-69.4 mole %. Thus, 160 g II in 200 ml  $\text{H}_2\text{O}$  was added to 100 g I in 200 ml  $\text{H}_2\text{O}$ , the whole mixed with 184 g 30%  $\text{SiO}_2$  soln, spray-dried, pelletized, calcined 4 hr at  $530^\circ$  in air, and milled to give a catalyst containing 25%  $\text{SiO}_2$ , which was used for manufacturing III to give  $\text{C}_3\text{H}_6$  conversion 84.3 mole % at III selectivity 63.4%.

IC C07C

CC 23-19 (Aliphatic Compounds)

IT 29075-42-1

RL: CAT (Catalyst use); USES (Uses)

(catalysts from telluric acid and, for oxidation of propylene)

IT 107-02-8P, preparation 107-13-1P, preparation


RL: IMF (Industrial manufacture); PREP (Preparation)  
(manufacture of, catalysts for)

IT 107-02-8P, preparation 107-13-1P, preparation

RL: IMF (Industrial manufacture); PREP (Preparation)  
(manufacture of, catalysts for)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 32 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1971:551381 HCAPLUS

DN 75:151381

TI Separation of acrylonitrile

IN Yamada, Yoshiteru; Nagai, Shigeki; Odan, Kyoji; Bando, Yasuo

PA Ube Industries, Ltd.

SO Jpn. Tokkyo Koho, 5 pp.

CODEN: JAXXAD

DT Patent

LA Japanese


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 46030817	B4	19710907	JP	19680924
AB	Acrylonitrile (I), MeCN, and acrolein were obtained from crude I (manufactured by ammoxidn. of propylene), vaporized, and treated with a mol. sieve composed of styrenedivinylbenzene copolymer to give pure I.				
IC	C07C; B01D				
CC	23 (Aliphatic Compounds)				
IT	107-13-1P, preparation				
	RL: PUR (Purification or recovery); PREP (Preparation) (purification of, acetonitrile and acrolein removal in)				
IT	75-05-8, uses and miscellaneous 107-02-8, uses and miscellaneous				

RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylonitrile)  
 IT 107-13-1P, preparation  
 RL: PUR (Purification or recovery); PREP (Preparation)  
 (purification of, acetonitrile and acrolein removal in)  
 RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



IT 107-02-8, uses and miscellaneous  
 RL: REM (Removal or disposal); PROC (Process)  
 (removal of, from acrylonitrile)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



L33 ANSWER 33 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1970:66350 HCAPLUS  
 DN 72:66350  
 TI 1,3,6-Octatriene  
 IN Feldman, Julian; Saffer, Bernard A.; Frampton, Orville D.  
 PA National Distillers and Chemical Corp.  
 SO U.S., 4 pp. Continuation-in-part of U.S. 3284529  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 3480685	A	19691125	US 1963-266514	19630320
PRAI	US 1963-266514	A	19630320		

AB Linear dimers, e.g. 1,3,6-octatriene (I), are prepared from open-chain conjugated diolefins, e.g., 1,3-butadiene (II); 2,6-octadiene and 2,4,6-octatriene are recovered as by-products. II is contacted with a Ni(O) carbonyl catalyst, e.g. (PhO)3PNi(CO)3, (Bu3P)2Ni(CO)2, (Ph3As)2Ni(CO)2, or [(PhO)3P]2Ni diacrolein, and a C3-10 aliphatic or alicyclic alc. as cocatalyst, at 70-160° 2-6 hr. to give a product containing 9-29% I. A polymerization inhibitor, e.g. p-tert-butylpyrocatechol, may be used.

IC C07C

INCL 260677000

CC 23 (Aliphatic Compounds)

IT 12120-60-4, Nickel, bis(acrylonitrile)bis(phosphorous acid)-, hexaphenyl ester 12567-69-0, Nickel, bis(acrolein)bis(phosphorous acid)-, cyclic diester with 2-(hydroxymethyl)-2-methyl-1,3-propanediol 12579-29-2, Nickel, bis(acrolein)bis(phosphorous acid)-, hexaphenyl ester 12580-97-1, Nickel, bis(cinnamonnitrile)bis(triphenylphosphine)- 13007-90-4 14653-44-2 15698-54-1 15709-52-1 18474-92-5 20658-46-2 28042-59-3

RL: CAT (Catalyst use); USES (Uses)

(catalysts, containing isopropyl alc. for octatriene manufacture)

IT 107-02-8DP, Acrolein, nickel complexes 107-13-1DP,


Acrylonitrile, nickel complexes 603-32-7DP, Arsine, triphenyl-, nickel complexes 603-35-0DP, Phosphine, triphenyl-, nickel complexes 603-36-1DP, Stibine, triphenyl-, nickel complexes 998-40-3DP, Phosphine, tributyl-, nickel complexes 1449-91-8DP, 1,3-Propanediol, 2-(hydroxymethyl)-2-methyl-, cyclic phosphite (1:1), nickel complexes 4360-47-8DP, Cinnamionitrile, nickel complexes

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

IT 107-02-8DP, Acrolein, nickel complexes 107-13-1DP,  
Acrylonitrile, nickel complexes  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 107-02-8 HCAPLUS

CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS

CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 34 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN

AN 1968:410107 HCAPLUS

DN 69:10107

TI Arsenophosphomolybdic acid catalysts for preparation of acrylonitrile and acrolein

IN Young, Howard S.

PA Eastman Kodak Co.

SO U.S., 3 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 3379652	A	19680423	US 1967-634772	19670428
PRAI	US 1967-634772	A	19670428		

AB Title acid (I) catalyst comps. useful in preparation of acrylonitrile (II) and acrolein are prepared from an As oxide, H<sub>3</sub>PMo<sub>12</sub>O<sub>40</sub> (III), and SiO<sub>2</sub>. Thus, a mixture containing dodecamolybdophosphoric acid hydrate 252, NH<sub>3</sub>-stabilized silica sol 825, As<sub>2</sub>O<sub>5</sub> oxide 1212, and H<sub>2</sub>O 250 g. was heated with stirring until it thickened to a yellow slurry, dried on a steam bath, and calcined 4 hrs. at 200° to give I. A mixture of propylene 214, air 1071, and steam 214 ml. was passed over 200 ml. I in a reactor at 475°, with a contact time 2.97 sec. Acrolein (11.6% yield) was obtained after 30 min. with some acrylic acid. II was prepared in 57.9% yield from propylene, O, and NH<sub>3</sub> by using I containing 5.2% As<sub>2</sub>O<sub>5</sub> and 41.6% III. Similarly prepared were methacrolein and methacrylic acid by reaction between isobutylene and O in the vapor phase in the presence of I.

INCL 252437000


CC 23 (Aliphatic Compounds)

IT 1303-28-2, Arsenic oxide (As<sub>2</sub>O<sub>5</sub>)

RL: CAT (Catalyst use); USES (Uses)

(catalysts from molybdophosphoric acid (H<sub>3</sub>PMo<sub>12</sub>O<sub>40</sub>) and

silica gel and, for oxidation of 2-methylpropene and propene)  
 IT 107-02-8P, preparation 107-13-1P, preparation  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (manufacture of)  
 IT 107-02-8P, preparation 107-13-1P, preparation  
 RL: IMF (Industrial manufacture); PREP (Preparation)  
 (manufacture of)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



L33 ANSWER 35 OF 35 HCAPLUS COPYRIGHT 2005 ACS on STN  
 AN 1967:28373 HCAPLUS  
 DN 66:28373  
 TI Linear octatrienes  
 IN Feldman, Julian; Saffer, Bernard A.; Frampton, Orville D.  
 PA National Distillers and Chemical Corp.  
 SO U.S., 7 pp.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 3284529		19661108	US	19621231
	DE 1468408			DE	
	GB 1061482			GB	

AB 1,3,7-(I) and 1,3,6-Octatrienes (II) were prepared by polymerizing 1,3-butadiene in the presence of 0.5-5.0% Ni (0) catalyst, in particular, bis(triphenyl phosphite)-nickel dicarbonyl (III), and 10-30% by weight m-cresol co-catalyst at 70-160°, separating the components of the product by fractional distillation, and separating I from the resulting binary system

of I and 4-vinyl-1-cyclohexene (IV) by thermal diffusion. Participation of the cresol in the reaction was demonstrated by the use of deuterated cresol. The linear octatrienes are useful in forming polymers. Thus, a 1-l. autoclave was charged with 5-10 g. ground CaC<sub>2</sub>, 72 g. m-cresol, 15 g. p-xylene, and 7 g. III purged with N, cooled, 419 g. chilled freshyl distilled 1,3-butadiene added, the chilled autoclave purged with N, closed, heated 20 hrs. with steam, cooled, vented, and the contents distilled through a Widmer column at atmospheric pressure to an overhead temperature of 175° to give 324 g. distillate and 177 g. residue. The distillate, as determined by vapor-phase chromatog, contained 34% I, 11% II, 9.2% and 34% 1,5-cyclooctadiene. By fractional distillation II and I could be separated from each other, but not from IV. IV was separated from I and II by liquid thermal diffusion in an apparatus described in detail. I b. 123°, d<sub>20</sub> 0.7640, n<sub>D</sub> 1.46446.

INCL 260677000

CC 23 (Aliphatic Compounds)  
 IT Phosphorous acid, triphenyl ester, nickel complexes  
 RL: CAT (Catalyst use); USES (Uses)  
 (catalysts from phenols and, for dimerization of 1,3-butadiene)  
 IT 107-02-8D, Acrolein, nickel complexes 107-13-1D,  
 Acrylonitrile, nickel complexes 603-35-0D, Phosphine, triphenyl-, nickel  
 complexes 764-42-1D, Fumaronitrile, nickel complexes 4360-47-8D,  
 Cinnamonnitrile, nickel complexes 12114-60-2 12120-60-4 13007-90-4  
 14653-44-2  
 RL: CAT (Catalyst use); USES (Uses)  
 (catalysts from phenols and, for dimerization of 1,3-butadiene)  
 IT 107-02-8D, Acrolein, nickel complexes 107-13-1D,  
 Acrylonitrile, nickel complexes  
 RL: CAT (Catalyst use); USES (Uses)  
 (catalysts from phenols and, for dimerization of 1,3-butadiene)  
 RN 107-02-8 HCAPLUS  
 CN 2-Propenal (9CI) (CA INDEX NAME)



RN 107-13-1 HCAPLUS  
 CN 2-Propenenitrile (9CI) (CA INDEX NAME)



=>